

FINLAND'S INFORMATIVE INVENTORY REPORT 2022

Air Pollutant Emissions 1980-2020
under the UNECE CLRTAP and the EU NECD

Part 1B - General

March 2022

FINNISH ENVIRONMENT INSTITUTE
Centre for Sustainable Consumption and Production
Environmental Management in Industry – Air Emissions Team

PART 1B - GENERAL

8.1 Recalculations and Improvements

8.1.1 Summary of recalculations, explanations and justifications

8.1.2 Impacts of recalculations to the 2022 submission

8.1.3 Improvements and Reviews

8.2 Projections

8.2.1 Projections for 2015, 2025 and 2030

8.2.2 Projections for Energy

8.2.3 Projections for Transport

8.2.4. Projections for IPPU and Waste

8.2.5 Projections for Agriculture

8.2.6 Emission reductions based on existing measures and measures that have been adopted in the legislation

8.2.7 NECD 2921 Review of Projections

8.3 Gridded Emissions and LPS

8.3.1 Gridded data

8.3.2 LPS data, sources, geographical coordinates and emissions

8.4 MEMO ITEMS

Overall description and methodologies

International aviation cruise (civil)

Domestic aviation cruise civil

International maritime navigation

Multilateral operations

Transport (fuel used)

Other not included in national total of the entire territory

Volcanoes

Forest fires

Other natural emissions

REFERENCES

8.1 RECALCULATIONS AND IMPROVEMENTS

Changes in chapter	
March 2022	KS

8.1.1 Summary of recalculations, explanations and justifications

Requested information

According to the Reporting Guidelines this chapter should include information relevant for assessment of compliance with each Protocol including a description of sources that were not included in the base year but have been added since as well as for sources that were included in the base year and are no longer applicable.

There was no obligation to document this information in the early years of reporting air pollutant emission inventories in the 1990's and the reporting guidelines have much developed since, therefore it has not been possible to present the requested information for the early years.

In the IIR sub-chapters "**Source specific recalculations**" of each Sector Chapter, information on annually conducted recalculations is presented for those years when the documentation item already existed.

Recalculations prior to the 2018 submission

The first full recalculation of the time series 1980-2016 was carried out to the submission in 2018. The recalculation of the energy sector time series from the 1990's was initiated in 2002, however, completed first to the 2018 submission. Due to the pending energy sector recalculations, it was not possible to fully recalculate interconnected data due to the complex structure of the inventory as explained in details in Chapter 2.3.2 in Part 1A – General of the IIR. However, individual emission figures and notation keys were corrected in the NFR tables when errors were found, in addition to sources where the activity data did not interfere with data reported by the plants. The ammonia emissions time-series was an exception, and was recalculated because sparse ammonia sources are related to data reported by the operators. The allocation of emissions under consistent reporting categories in the time series was not realized until the full recalculation to the 13 April 2018 resubmission. No impact assessments of the partial recalculations until 2018 were performed due to resource limitations and the fact that the impact on the non-recalculated time series would anyway be highly uncertain.

Resubmissions in 2018-2022

15 Feb 2018 Finland submitted the old time series for the years 1990-2015 and new data for 2016. This was because the energy sector data was not finalized by the deadline of the NFR tables.

15 Mar 2018 Finland submitted the first recalculated time series, however, it would not have been mature for submission due to lack of checks that could not be done in the window between the late finalization of the energy sector data and the 4 weeks time frame for resubmissions.

13 Apr 2018 Finland submitted a recalculated time series that had undergone several QA/QC procedures, however, still having remaining reallocation issues. Due to the UNECE CLRTAP S3 Review and the EU NECD Technical Review, both in June 2018, the data needed to be available.

15 Feb 2019 Finland submitted the recalculated time series which included further harmonized

emissions allocations in the time series, however, also some errors were discovered after the submission deadline

15 Mar 2019 Finland submitted additional corrections to the submission of 15 Feb 2019.









13 Mar 2020 Finland submitted corrections to the submission of 13 Feb 2020 (errors and omissions) due to time constraint caused by unexpected data flows: (1) renewal of the contents of the YLVA database with deletion of technical details used in the energy sector inventory and pre-scheduled initiation of the new energy sector calculation model, (2) errors identified in the agriculture sector calculation model formulas, (3) omission of recalculated values from the submission (agriculture HCB), (4) missing values not captured into the submission 13 February 2020. The impacts of the recalculations are presented in Chapter 8.2.









15 Mar 2022 Finland submitted corrections due to errors in NH₃ from 5B2 and some AD changes. In addition, the 2019 adjustment values for 2019 were accidentally included for the year 2020 and were removed.

8.1.2 Impacts of recalculations in the 2022 submission

Recalculations to the 2022 submission are explained in detail in the respective IIR chapters (Parts 2-6 of the IIR). Below is an overview of the impacts of the recalculations to the 2022 submission in Figure 8.1 and Table 8.1. Most of the recalculations are due to update of statistical data and in some cases application of new emission factors, the largest changes being due to:

- *Heavy metal emissions 1997 are corrected due to addition of missing emissions from petroleum coke at refineries*
- *2019 PCDD/F emissions are revised due to corrected reporting of a metal industry plant*
- *PAH-4 and particle emissions along the time series are revised due to corrections in Statistics Finland's wood use statistics*
- *PCB 2017-2019 are revised due to correction of activity data*
- *NH₃ whole timeseries: see IIR Part 4 Agriculture, chapters Source-specific recalculations including changes made in response to the review process.*








Figure 8.1 Impact of recalculations in the 2022 submission compared to the 2021 submission

Table 8.1 Impact of recalculations in the 2022 submission compared to the 2021 submission

NOx															
	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994
15 Feb 2021	307.182	286.884	282.262	273.176	269.054	286.991	288.711	300.116	303.080	309.551	306.359	303.483	287.916	293.134	293.859
4 March 2022	307.310	287.732	282.580	273.461	269.288	287.133	288.909	300.330	303.293	309.780	306.550	303.652	288.070	293.281	294.000
Difference	0.127	0.848	0.318	0.284	0.234	0.143	0.198	0.214	0.213	0.229	0.192	0.169	0.154	0.147	0.141
%	0.0	0.3	0.1	0.1	0.1	0.0	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0
	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
15 Feb 2021	273.114	277.454	271.585	257.440	252.746	241.062	244.344	242.180	248.635	237.073	208.102	223.729	210.841	193.525	176.440
15 March 2020	273.027	277.389	271.555	257.426	252.773	241.124	244.475	242.313	248.767	237.202	208.230	223.853	210.960	193.626	176.458
Difference	-0.087	-0.065	-0.030	-0.014	0.027	0.062	0.131	0.133	0.132	0.129	0.128	0.124	0.119	0.101	0.018
%	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0
	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019					
15 Feb 2021	187.294	171.303	161.521	158.398	150.714	138.782	134.600	130.2220426	126.721	119.817					
4 March 2022	187.211	171.310	161.436	158.508	150.832	138.933	134.633	130.379	127.089	119.901					
Difference	-0.083	0.007	-0.085	0.111	0.118	0.151	0.032	0.157	0.368	0.084					
%	0.0	0.0	-0.1	0.1	0.1	0.1	0.0	0.1	0.3	0.1					
SOx															
	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994
15 Feb 2021	584.393	534.327	484.234	372.240	368.242	382.271	331.171	328.281	302.271	244.256	248.820	205.522	156.202	137.604	122.681
4 March 2022	585.029	534.931	484.785	372.765	368.704	382.708	331.549	328.697	302.705	244.651	248.820	205.522	156.202	137.604	122.681
Difference	0.635	0.604	0.552	0.524	0.463	0.437	0.378	0.416	0.433	0.395	0.000	0.000	0.000	0.000	0.000
%	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.0	0.0	0.0	0.0	0.0
	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
15 Feb 2021	104.555	109.085	100.679	93.411	91.842	81.812	95.775	90.387	101.105	83.529	69.5506	82.91414	81.16953	66.83803	58.91219
4 March 2022	104.542	109.073	100.669	93.402	91.836	81.808	95.775	90.387	101.105	83.529	69.55065	82.91414	81.16953	66.83712	58.90614
Difference	-0.013	-0.012	-0.010	-0.009	-0.006	-0.005	0.000	0.000	0.000	0.000	0.000	0.000	0.000	-0.001	-0.006
%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019					
15 Feb 2021	66.10813	60.22403	49.96141	47.61231	44.25089	40.82686	39.81346	35.02135	33.12463	28.93676					
4 March 2022	66.0958	60.2174	49.94922	47.6123	44.25118	40.82905	39.80869	35.02096	33.12461	29.59191					
Difference	-0.012	-0.007	-0.012	0.000	0.000	0.002	-0.005	0.000	0.000	0.655					
%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.2					

NH3

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994
15 Feb 2021	35.92833	36.33223	36.79751	36.94373	36.86605	36.79542	36.39727	35.5807	34.81443	33.13697	34.73792	33.19021	31.8574	32.28547	33.2658
4 March 2022	36.93201	37.35076	37.80853	37.93955	37.87626	37.78855	37.37521	36.52977	35.73182	34.01788	35.62384	34.03729	32.68716	33.13695	34.11939
Difference	1.004	1.019	1.011	0.996	1.010	0.993	0.978	0.949	0.917	0.881	0.886	0.847	0.830	0.851	0.854
%	2.7	2.7	2.7	2.6	2.7	2.6	2.6	2.6	2.6	2.6	2.5	2.5	2.5	2.6	2.5
	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
15 Feb 2021	33.42462	33.42462	35.98686	35.25875	37.9425	34.815	35.012	35.854	36.809	37.270	37.996	37.028	36.600	35.757	36.265
4 March 2022	34.13693	34.13693	36.74254	36.02992	38.7415	35.656	35.904	36.777	37.735	38.203	38.940	37.978	37.532	36.683	37.186
Difference	0.712	0.712	0.756	0.771	0.799	0.841	0.892	0.923	0.926	0.933	0.943	0.950	0.931	0.926	0.921
%	2.1	2.1	2.1	2.1	2.1	2.4	2.5	2.5	2.5	2.4	2.4	2.5	2.5	2.5	2.5
	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019					
15 Feb 2021	36.596	35.345	35.247	34.592	35.150	33.533	33.192	32.331	32.121	31.593					
4 March 2022	37.560	36.324	36.240	35.565	36.089	34.367	34.020	33.142	32.924	32.181					
Difference	0.964	0.979	0.993	0.973	0.938	0.835	0.828	0.811	0.802	0.588					
%	2.6	2.7	2.7	2.7	2.6	2.4	2.4	2.4	2.4	1.8					

NM VOC

	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999							
15 Feb 2021	239.7604	233.1867	233.0104	223.2739	217.2101	211.0293	210.1266	203.166	195.6615	195.5312	190.6597	184.2842							
4 March 2022	240.04	233.44	235.51	225.78	219.71	213.53	212.71	204.50	197.25	197.30	192.60	186.34							
Difference	0.281	0.254	2.504	2.503	2.496	2.504	2.588	1.336	1.592	1.764	1.943	2.053							
%	0.1	0.1	1.1	1.1	1.1	1.2	1.2	0.7	0.8	0.9	1.0	1.1							
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018
15 Feb 2021	177.9705	175.0953	166.1791	161.9135	157.3166	146.2027	140.1557	136.4293	120.3039	111.3182	113.4094	103.9307	101.5742	96.24821	93.66584	88.89524	89.83256	87.35063	85.43574
4 March 2022	178.98	176.57	167.96	163.51	158.56	147.66	141.58	137.93	121.78	113.09	113.77	105.20	101.73	97.44	95.11	89.96	90.45	88.36	86.48
Difference	1.008	1.475	1.783	1.596	1.246	1.460	1.422	1.496	1.477	1.774	0.357	1.269	0.160	1.188	1.440	1.069	0.614	1.008	1.045
%	0.6	0.8	1.1	1.0	0.8	1.0	1.0	1.1	1.2	1.6	0.3	1.2	0.2	1.2	1.5	1.2	0.7	1.1	1.2

CO

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999									
15 Feb 2021	769.7768	741.224	720.5285	705.6586	692.1802	675.3056	669.8025	663.097	657.5392	639.4846									
4 March 2022	764.4565	735.8418	715.1348	700.2468	686.5365	661.765	656.8048	651.3897	646.2599	629.839									
Difference	-5.320	-5.382	-5.394	-5.412	-5.644	-13.541	-12.998	-11.707	-11.279	-9.646									
%	-0.7	-0.7	-0.8	-0.8	-0.8	-2.0	-2.0	-1.8	-1.7	-1.5									
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018
15 Feb 2021	601.1868	601.362	582.8371	561.8072	547.4071	523.8428	503.5113	490.4429	456.7624	437.0613	453.841	411.7636	409.3276	389.1865	382.9343	359.5816	370.7473	358.9465	350.0619
4 March 2022	593.7654	596.0474	577.4972	556.3416	541.87	519.4174	499.2473	486.0342	452.0152	428.96	446.0043	407.361	401.5643	388.655	382.5706	359.4196	366.2629	357.1588	348.6837
Difference	-7.421	-5.315	-5.340	-5.466	-5.537	-4.425	-4.264	-4.409	-4.747	-8.101	-7.837	-4.403	-7.763	-0.531	-0.364	-0.162	-4.484	-1.788	-1.378
%	-1.2	-0.9	-0.9	-1.0	-1.0	-0.9	-0.9	-0.9	-1.1	-1.9	-1.8	-1.1	-1.9	-0.1	-0.1	0.0	-1.2	-0.5	-0.4

TSP

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999									
15 Feb 2021	98.90518	86.22526	78.83221	73.58427	73.72423	68.43487	65.81213	65.4201	59.39055	61.11393									
4 March 2022	98.73523	86.06132	78.6656	73.40887	73.58643	67.75057	65.16779	64.84385	58.85361	60.67784									
Difference	-0.170	-0.164	-0.167	-0.175	-0.138	-0.684	-0.644	-0.576	-0.537	-0.436									
%	1.0	1.2	1.3	1.3	1.8	0.9	1.2	1.2	1.3	1.4									
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018
15 Feb 2021	56.8545	58.45997	59.63142	60.67307	59.09387	57.2283	59.00351	55.60479	52.94356	52.09021	54.36649	51.29598	49.07064	49.00717	48.37681	45.12334	47.39466	45.15251	45.05578
4 March 2022	56.41023	58.15065	59.32278	60.35906	58.77509	57.07477	58.86147	55.46177	52.78474	51.73366	53.665	50.83512	48.39084	48.7815	48.15705	44.79637	46.78251	44.7083	44.63484
Difference	-0.444	-0.309	-0.309	-0.314	-0.319	-0.154	-0.142	-0.143	-0.159	-0.357	-0.701	-0.461	-0.680	-0.226	-0.220	-0.327	-0.612	-0.444	-0.421
%	-0.8	-0.5	-0.5	-0.5	-0.5	-0.3	-0.2	-0.3	-0.3	-0.7	-1.3	-0.9	-1.4	-0.5	-0.5	-0.7	-1.3	-1.0	-0.9

PM10

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999										
15 Feb 2021	74.17929	66.8966	60.77715	56.54723	55.8371	51.66347	50.2302	49.68909	45.79967	46.08663										
4 March 2022	74.01701	66.74289	60.62126	56.38256	55.7105	51.01051	49.61543	49.12751	45.28756	45.67188										
Difference	-0.162	-0.154	-0.156	-0.165	-0.127	-0.653	-0.615	-0.562	-0.512	-0.415										
%	-0.2	-0.2	-0.3	-0.3	-0.2	-1.3	-1.2	-1.1	-1.1	-0.9										
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
15 Feb 2021	42.93129	43.91933	44.34985	45.32758	44.13141	42.32309	43.14736	40.98366	38.56066	37.47466	39.05181	36.02412	34.89423	34.33282	33.81448	31.37474	32.57913	30.98986	31.10069	30.03414
4 March 2022	42.50976	43.6275	44.05862	45.03175	43.83072	42.18123	43.01723	40.85351	38.41577	37.13906	38.38808	35.59344	34.2546	34.12937	33.6176	31.07565	32.00267	30.57583	30.69854	29.61065
Difference	-0.422	-0.292	-0.291	-0.296	-0.301	-0.142	-0.130	-0.130	-0.145	-0.336	-0.664	-0.431	-0.640	-0.203	-0.197	-0.299	-0.576	-0.414	-0.402	-0.423
%	-1.0	-0.7	-0.7	-0.7	-0.7	-0.3	-0.3	-0.3	-0.4	-0.9	-1.7	-1.2	-1.9	-0.6	-0.6	-1.0	-1.8	-1.4	-1.3	-1.4

PM2.5

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999										
15 Feb 2021	47.39727	42.77847	38.96607	35.59987	35.02147	32.50242	31.64716	30.95806	28.64718	28.7607										
4 March 2022	47.24757	42.62757	38.81473	35.44752	34.86445	31.86712	31.05391	30.43217	28.15488	28.36289										
Difference	-0.150	-0.151	-0.151	-0.152	-0.157	-0.635	-0.593	-0.526	-0.492	-0.398										
%	-0.3	-0.4	-0.4	-0.4	-0.5	-2.0	-1.9	-1.7	-1.7	-1.4										
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
15 Feb 2021	26.32582	26.99546	27.13458	27.43044	26.85738	26.02572	25.70645	24.54568	22.81868	22.21083	23.79563	20.87371	20.80342	19.69909	19.38443	17.6904	18.60342	17.85298	17.8082	16.62195
4 March 2022	25.92067	26.71772	26.85657	27.14763	26.57184	25.89589	25.5819	24.42243	22.68034	21.88654	23.15385	20.45866	20.18036	19.50549	19.19845	17.40377	18.04568	17.45406	17.41977	16.20936
Difference	-0.405	-0.278	-0.278	-0.283	-0.286	-0.130	-0.125	-0.123	-0.138	-0.324	-0.642	-0.415	-0.623	-0.194	-0.186	-0.287	-0.558	-0.399	-0.388	-0.413
%	-1.6	-1.0	-1.0	-1.0	-1.1	-0.5	-0.5	-0.5	-0.6	-1.5	-2.8	-2.0	-3.1	-1.0	-1.0	-1.6	-3.1	-2.3	-2.2	-2.5

BC

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999										
15 Feb 2021	10.10017	9.601714	9.151792	8.718234	8.548016	7.956742	7.668376	7.42125	7.246812	6.961374										
4 March 2022	10.13737	9.639547	9.189432	8.756007	8.587378	7.846707	7.577087	7.35197	7.190166	6.929695										
Difference	0.037	0.038	0.038	0.038	0.039	-0.110	-0.091	-0.069	-0.057	-0.032										
%	0.4	0.4	0.4	0.4	0.5	-1.4	-1.2	-0.9	-0.8	-0.5										
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
15 Feb 2021	6.500459	6.673922	6.580453	6.440257	6.295144	5.899511	5.527889	5.512307	5.314556	5.314556	5.599238	4.873316	5.088504	4.632816	4.527093	4.152249	4.351033	4.12788	3.979532	3.847619
4 March 2022	6.489547	6.715374	6.622459	6.483719	6.33936	5.950592	5.576581	5.563682	5.356405	5.356405	5.465733	4.800154	4.959388	4.627451	4.524986	4.130808	4.256194	4.078406	3.92944	3.798568
Difference	-0.011	0.041	0.042	0.043	0.044	0.051	0.049	0.051	0.042	0.042	-0.134	-0.073	-0.129	-0.005	-0.002	-0.021	-0.095	-0.049	-0.050	-0.049
%	-0.2	0.6	0.6	0.7	0.7	0.9	0.9	0.9	0.8	0.8	-2.4	-1.5	-2.6	-0.1	0.0	-0.5	-2.2	-1.2	-1.3	-1.3

Pb

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999										
15 Feb 2021	321.4349	237.157	165.1449	105.1439	73.96598	72.71721	49.24703	31.87141	37.26168	33.68885										
4 March 2022	321.4349	237.157	165.1449	105.1439	73.96598	72.69021	49.22303	34.3328	37.24368	33.67585										
Difference	0.000	0.000	0.000	0.000	0.000	-0.027	-0.024	2.461	-0.018	-0.013										
%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	7.2	0.0	0.0										
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
15 Feb 2021	30.67128	29.79993	30.73762	24.92638	26.49538	21.4896	24.86589	21.85402	19.82316	16.77146	20.39779	19.24543	16.36296	16.01024	16.64614	14.69011	15.7196	15.65635	15.42797	13.22047
4 March 2022	30.66228	29.79993	30.73762	24.92638	26.49538	21.4897	24.86589	21.85402	19.82133	16.75936	20.37314	19.23217	16.33858	16.01021	16.64672	14.69449	15.71005	15.64938	15.42887	13.21338
Difference	-0.009	0.000	0.000	0.000	0.000	0.000	0.000	0.000	-0.002	-0.012	-0.025	-0.013	-0.024	0.000	0.001	0.004	-0.010	-0.007	0.001	-0.007
%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	-0.1	-0.1	-0.1	-0.1	0.0	0.0	0.0	-0.1	0.0	0.0	-0.1

Cd																				
	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999										
15 Feb 2021	6.685065	3.810535	3.326164	3.386844	2.71969	2.134925	1.917333	1.539125	1.699365	1.538388										
4 March 2022	6.685065	3.810535	3.326164	3.386844	2.71969	2.126825	1.910133	1.589663	1.693965	1.534488										
Difference	0.000	0.000	0.000	0.000	0.000	-0.008	-0.007	0.051	-0.005	-0.004										
%	0.0	0.0	0.0	0.0	0.0	-0.4	-0.4	3.2	-0.3	-0.3										
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
15 Feb 2021	1.414031	1.745094	1.374596	1.327277	1.615047	1.450951	1.426068	1.266091	1.206573	1.156997	1.290374	1.222736	1.176966	1.079379	0.918999	0.88799	0.942545	0.95496	0.880947	0.793907
4 March 2022	1.411331	1.745094	1.374596	1.327277	1.615047	1.450981	1.426068	1.266091	1.206024	1.153367	1.282979	1.218758	1.169652	1.07937	0.919173	0.889304	0.93968	0.95473	0.880934	0.788896
Difference	-0.003	0.000	0.000	0.000	0.000	0.000	0.000	0.000	-0.001	-0.004	-0.007	-0.004	-0.007	0.000	0.000	0.001	-0.003	0.000	0.000	-0.005
%	-0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	-0.3	-0.6	-0.3	-0.6	0.0	0.0	0.1	-0.3	0.0	0.0	-0.6
Hg																				
	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999										
15 Feb 2021	1.086065	0.929457	0.890854	0.765221	0.808063	0.80189	0.866198	0.802507	0.693267	0.636198										
4 March 2022	1.086065	0.929457	0.890854	0.765221	0.808063	0.80054	0.864998	0.813129	0.692367	0.635548										
Difference	0.000	0.000	0.000	0.000	0.000	-0.001	-0.001	0.011	-0.001	-0.001										
%	0.2	0.3	0.3	0.3	0.5	0.3	0.4	0.4	0.5	0.6										
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
15 Feb 2021	0.603664	0.684325	0.671836	0.818907	0.753327	0.891222	1.034303	0.872876	0.820613	0.756448	0.88855	0.88855	0.745813	0.761606	0.706815	0.62093	0.591037	0.581787	0.679154	0.586585
4 March 2022	0.603214	0.684325	0.671836	0.818907	0.753327	0.891226	1.034303	0.872875	0.820521	0.755842	0.887317	0.887317	0.744593	0.761603	0.706842	0.621147	0.590558	0.581745	0.67915	0.586585
Difference	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	-0.001	-0.001	-0.001	-0.001	0.000	0.000	0.000	0.000	0.000	0.000	0.000
%	-0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	-0.1	-0.1	-0.1	-0.2	0.0	0.0	0.0	-0.1	0.0	0.0	0.0
As																				
	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999										
15 Feb 2021	34.81587	23.85912	18.04396	16.38553	10.99594	5.224612	8.500685	13.45645	13.98414	5.207524										
4 March 2022	34.81587	23.85912	18.04396	16.38553	10.99594	5.221912	8.498285	13.86278	13.98234	5.206224										
Difference	0.000	0.000	0.000	0.000	0.000	-0.003	-0.002	0.406	-0.002	-0.001										
%	0.0	0.0	0.0	0.0	0.0	-0.1	0.0	2.9	0.0	0.0										
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
15 Feb 2021	4.362116	5.121256	3.990792	3.668156	4.112212	2.991875	2.945346	2.959251	3.135908	2.893816	3.401971	2.982599	2.644915	2.755253	2.708919	2.456251	2.579076	2.446004	2.414967	2.064919
4 March 2022	4.361216	5.121256	3.990792	3.668156	4.112212	2.991885	2.945346	2.959251	3.135725	2.892606	3.399506	2.981273	2.642477	2.75525	2.708977	2.456689	2.578121	2.445923	2.408086	2.05761
Difference	-0.001	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	-0.001	-0.002	-0.001	-0.002	0.000	0.000	0.000	-0.001	0.000	-0.007	-0.007
%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	-0.1	0.0	-0.1	0.0	0.0	0.0	0.0	0.0	-0.3	-0.4

Cr																				
	13.980																			
	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999										
15 Feb 2021	47.66979	59.68513	47.62388	37.7212	40.66784	35.92104	32.83785	29.55647	30.13295	31.2715										
4 March 2022	47.66979	59.68513	47.62388	37.7212	40.66784	35.82654	32.75385	32.62747	30.06995	31.226										
Difference	0.000	0.000	0.000	0.000	0.000	-0.095	-0.084	3.071	-0.063	-0.046										
%	0.0	0.0	0.0	0.0	0.0	-0.3	-0.3	9.4	-0.2	-0.1										
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	
15 Feb 2021	28.59477	25.69396	39.07483	29.45505	26.03354	20.00778	24.92144	28.82415	26.51467	16.7039	25.97075	17.41695	18.64387	18.43414	23.06255	16.70088	18.2056	16.6541	15.34341	14.28566
4 March 2022	28.56327	25.69396	39.07483	29.45505	26.03354	20.00813	24.92143	28.82415	26.50826	16.66155	25.88447	17.37054	18.55854	18.43403	23.06458	16.71621	18.17217	16.65135	15.34321	14.28571
Difference	-0.032	0.000	0.000	0.000	0.000	0.000	0.000	0.000	-0.006	-0.042	-0.086	-0.046	-0.085	0.000	0.002	0.015	-0.033	-0.003	0.000	0.000
%	-0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	-0.3	-0.3	-0.3	-0.5	0.0	0.0	0.1	-0.2	0.0	0.0	0.0
Cu																				
	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999										
15 Feb 2021	156.8926	148.8692	124.2826	111.5387	106.1926	116.5008	109.5383	127.9255	84.48406	68.13224										
4 March 2022	156.8926	148.8692	124.2826	111.5387	106.1926	116.4846	109.5239	129.0128	84.47326	68.12444										
Difference	0.000	0.000	0.000	0.000	0.000	-0.016	-0.014	1.087	-0.011	-0.008										
%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	0.0	0.0										
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	
15 Feb 2021	65.33805	66.23947	69.27718	61.89834	59.98121	58.05136	58.6445	44.14587	42.04309	40.42117	41.97953	42.4087	41.21558	42.37797	43.07401	40.74091	41.64717	40.63312	40.15571	40.17103
4 March 2022	65.33265	66.23947	69.27718	61.89834	59.98121	58.05142	58.6445	44.14587	42.04199	40.41391	41.96473	42.40074	41.20095	42.37795	43.07435	40.74353	41.64144	40.63435	40.06925	40.07676
Difference	-0.005	0.000	0.000	0.000	0.000	0.000	0.000	0.000	-0.001	-0.007	-0.015	-0.008	-0.015	0.000	0.000	0.003	-0.006	0.001	-0.086	-0.094
%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	-0.2	-0.2
Ni																				
	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999										
15 Feb 2021	78.43879	60.8846	52.20383	45.62521	44.61004	46.66409	36.86966	38.17576	33.6169	37.14945										
4 March 2022	78.43879	60.8846	52.20383	45.62521	44.61004	46.58309	36.79766	38.11576	33.5629	37.11045										
Difference	0.000	0.000	0.000	0.000	0.000	-0.081	-0.072	-0.060	-0.054	-0.039										
%	0.0	0.0	0.0	0.0	0.0	-0.2	-0.2	-0.2	-0.2	-0.1										
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
15 Feb 2021	34.59879	32.06462	37.91689	35.27905	30.56539	26.06078	27.82513	24.97175	22.35324	20.51056	22.85559	19.94886	18.94263	17.14711	17.31668	16.09092	15.74508	14.78368	14.14866	11.54071
4 March 2022	34.57179	32.06462	37.91689	35.27905	30.56539	26.06108	27.82513	24.97175	22.34775	20.47426	22.78164	19.90908	18.86949	17.14702	17.31842	16.10406	15.71643	14.78135	14.1485	11.53827
Difference	-0.027	0.000	0.000	0.000	0.000	0.000	0.000	0.000	-0.005	-0.036	-0.074	-0.040	-0.073	0.000	0.002	0.013	-0.029	-0.002	0.000	-0.002
%	-0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	-0.2	-0.3	-0.2	-0.4	0.0	0.0	0.1	-0.2	0.0	0.0	0.0

Zn																				
	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999										
15 Feb 2021	682.8393	472.7985	374.2281	349.3833	405.625	404.8758	271.4966	148.3986	152.1169	142.3637										
4 March 2022	682.8393	472.7985	374.2281	349.3833	405.625	402.9858	269.8166	151.7101	150.8569	141.4537										
Difference	0.000	0.000	0.000	0.000	0.000	-1.890	-1.680	3.312	-1.260	-0.910										
%	0.0	0.0	0.0	0.0	0.0	-0.5	-0.6	2.2	-0.8	-0.6										
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	
15 Feb 2021	128.261	131.3268	146.8727	127.0915	124.7922	119.3004	119.065	108.3952	116.819	116.6799	130.307	125.0412	128.5161	124.4094	131.8971	118.3094	128.0142	120.0158	118.6562	130.305
4 March 2022	127.631	131.3268	146.8727	127.0915	124.7922	119.3074	119.065	108.3952	116.6909	115.8329	128.5815	124.113	126.8095	124.4073	131.9377	118.616	127.3457	119.9608	118.4899	130.0144
Difference	-0.630	0.000	0.000	0.000	0.000	0.007	0.000	0.000	-0.128	-0.847	-1.726	-0.928	-1.707	-0.002	0.041	0.307	-0.669	-0.055	-0.166	-0.291
%	-0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	-0.1	-0.7	-1.3	-0.7	-1.3	0.0	0.0	0.3	-0.5	0.0	-0.1	-0.2
PCDD/F																				
	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999										
15 Feb 2021	17.62568	18.4154	17.35114	17.81984	18.16254	18.60414	16.98334	17.05507	17.1707	17.2528										
4 March 2022	17.54643	18.3539	17.30663	17.74441	18.08574	18.47767	16.85932	16.93457	17.04031	17.13511										
Difference	-0.079	-0.061	-0.045	-0.075	-0.077	-0.126	-0.124	-0.121	-0.130	-0.118										
%	-0.5	-0.3	-0.3	-0.4	-0.4	-0.7	-0.7	-0.7	-0.8	-0.7										
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
15 Feb 2021	17.68129	14.88651	15.12416	13.70747	13.85224	13.55029	14.05034	13.81586	13.83801	12.10372	15.91427	13.6386	14.5824	14.83329	15.64955	13.93468	14.94105	12.50627	13.41547	12.13155
4 March 2022	17.58616	14.79949	15.01973	13.63971	13.78918	13.48914	13.96508	13.74215	13.76279	12.01984	15.80232	13.56094	14.479	14.78034	15.59281	13.88096	14.87612	12.45187	13.36565	9.505828
Difference	-0.095	-0.087	-0.104	-0.068	-0.063	-0.061	-0.085	-0.074	-0.075	-0.084	-0.112	-0.078	-0.103	-0.053	-0.057	-0.054	-0.065	-0.054	-0.050	-2.626
%	-0.5	-0.6	-0.7	-0.5	-0.5	-0.5	-0.6	-0.5	-0.5	-0.7	-0.7	-0.6	-0.7	-0.4	-0.4	-0.4	-0.4	-0.4	-0.4	-27.6
PAH-4																				
	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999										
15 Feb 2021	18.96773	19.12122	19.16386	19.434	20.26539	19.45657	20.62231	20.57398	21.21191	20.57914										
4 March 2022	18.42696	18.57417	18.61563	18.88394	19.69175	17.7846	19.04068	19.16366	19.89207	19.46476										
Difference	-0.541	-0.547	-0.548	-0.550	-0.574	-1.672	-1.582	-1.410	-1.320	-1.114										
%	-2.9	-2.9	-2.9	-2.9	-2.9	-9.4	-8.3	-7.4	-6.6	-5.7										
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
15 Feb 2021	18.56488	20.60111	20.76537	21.31431	21.59015	22.22801	21.29508	22.22946	21.98663	23.47024	26.14006	22.28911	24.65095	22.52824	22.8106	21.37994	23.82096	23.03498	22.62306	22.30933
4 March 2022	18.07227	20.46292	20.6255	21.1691	21.44081	22.24609	21.30783	22.23534	21.91236	22.98243	25.55631	22.09924	24.05102	22.86818	23.17876	21.99621	23.91921	23.46506	23.07536	22.75325
Difference	-0.493	-0.138	-0.140	-0.145	-0.149	0.018	0.013	0.006	-0.074	-0.488	-0.584	-0.190	-0.600	0.340	0.368	0.616	0.098	0.430	0.452	0.444
%	-2.7	-0.7	-0.7	-0.7	-0.7	0.1	0.1	0.0	-0.3	-2.1	-2.3	-0.9	-2.5	1.5	1.6	2.8	0.4	1.8	2.0	2.0

HCB

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
15 Feb 2021	35.67656	35.56662	36.36116	36.37946	36.05407	35.58531	38.01543	38.46767	38.38615	38.1324
4 March 2022	35.67656	35.56662	36.36116	36.37946	36.05407	35.57181	38.00343	38.45767	38.37715	38.1259
Difference	0.000	0.000	0.000	0.000	0.000	-0.014	-0.012	-0.010	-0.009	-0.007
%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0

	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
15 Feb 2021	38.60062	18.18638	12.02426	9.912359	26.03888	32.37445	36.12212	38.18197	18.69147	26.57396	8.761446	26.11987	9.475741	17.30678	21.62263	16.20337	59.70087	33.40531	31.88326	22.63715
4 March 2022	38.59612	18.18638	12.02426	9.912358	26.03888	32.3745	36.12212	38.18197	18.69055	26.5679	8.749111	26.11322	9.463535	17.30675	21.6229	16.20554	59.69607	33.40489	31.88323	22.62692
Difference	-0.005	0.000	0.000	0.000	0.000	0.000	0.000	0.000	-0.001	-0.006	-0.012	-0.007	-0.012	0.000	0.000	0.002	-0.005	0.000	0.000	-0.010
%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	-0.1	0.0	-0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0

PCB

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
15 Feb 2021	28.8501	25.01989	26.12092	28.26471	28.91709	29.08038	27.8575	29.88556	31.71766	30.54547
4 March 2022	28.55767	24.78965	25.92362	28.00707	28.65386	28.67167	27.4561	29.48146	31.31546	30.16487
Difference	-0.292	-0.230	-0.197	-0.258	-0.263	-0.409	-0.401	-0.404	-0.402	-0.381
%	-1.0	-0.9	-0.8	-0.9	-0.9	-1.4	-1.5	-1.4	-1.3	-1.3

	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
15 Feb 2021	30.37051	29.46048	29.06863	30.25188	31.41095	31.4672	31.79738	32.24676	31.23548	21.29457	27.97779	27.61778	24.65171	23.49761	24.58694	24.17886	25.72329	25.7696	26.2604	22.7799
4 March 2022	30.03747	29.15423	28.72973	29.96413	31.126	31.19003	31.46568	31.93678	30.93202	20.96989	27.56623	27.29002	24.25722	23.24922	24.34253	23.95674	25.41757	22.68757	23.25623	19.7592
Difference	-0.333	-0.306	-0.339	-0.288	-0.285	-0.277	-0.332	-0.310	-0.303	-0.325	-0.412	-0.328	-0.394	-0.248	-0.244	-0.222	-0.306	-3.082	-3.004	-3.021
%	-1.1	-1.1	-1.2	-1.0	-0.9	-0.9	-1.1	-1.0	-1.0	-1.5	-1.5	-1.2	-1.6	-1.1	-1.0	-0.9	-1.2	-13.6	-12.9	-15.3

8.2 Improvements and Reviews

Inventory improvement programme at Finnish Environment Institute

Identification of further development needs in the Finnish air pollutant inventory is carried out on a continuous basis according to annual work programmes although larger scale improvements are possible only when the necessary resources for the improvement projects are available.

In the past years the inventory improvement programme was strongly linked with the national emission data production methods provided to the operators in their reporting to emission registers such as the E-PRTR. Finnish Environment Institute maintains information on emission estimation methodologies and emission factors on a website (http://www.ymparisto.fi/fi-FI/Asiointi_luvat_ja_ymparistovaikutusten_arviointi/Luvat_ilmoitukset_ja_rekisterointi/Paastotiedon_ilmoittaminen_paastorekistereihin_PRTR) (in Finnish). These methods should be applied in the E-PRTR reporting by the plant operators whenever no plant specific data is available. This procedure has been developed to ensure consistency between the data reported by the plants and the emission inventory.

The programme has thus far included studies in the energy production sector (boilers >50 MW), industrial processes (pulp and paper, iron and steel), agriculture and waste sectors and resulted in updating or developing of several emission factors. The studies involve also examination of the applicability of the default methods presented in the Guidebook for the national conditions.

National emission factors are derived from data reported by the plants when these are based on site-specific measurements and other site-specific data. In the later years, the obligation to use the latest version of the Guidebook emission factors has been more dominant. After the full recalculation of the time series emphasis will be given to check and further develop national emission factors based on data reported by the plants and replace the Guidebook EFs with these where feasible.

The results of the uncertainty analysis are used to prioritise the improvements.

Review, Improvement and Harmonization of the Nordic Air Emission Inventories in the Nordic Air Emission Experts Group

Changes in chapter	
March 2021	KS

Since 2004 the Nordic countries (Denmark, Finland, Iceland, Norway and Sweden) have carried out several projects on reviewing, improving and harmonizing the national air pollutant emission inventories. The work has been funded by the Nordic Council of Ministers. The target of the cooperation is to share knowledge and resources and to increase the quality of the Nordic CLRTAP air emission inventories with respect to accuracy, comparability, transparency and completeness. Until now, POP, NMVOC, particle and partly also heavy metal emission inventories in the Nordic countries have improved. Several improvements to the national inventories have been made in all Nordic countries due to the results of the work, for instance in NMVOC and particle emission inventories.

In addition to the overall review (2004), the following specific sectors have been under work:

- particulate emissions from small scale wood combustion and road transport (2006)
- emissions from the use of products (2006-2011)
- NMVOC inventories from the domestic product use sector (2010)
- SLCP emissions (2014-2017)
- POP and heavy metals from all sectors (2016-2018).
- POP and heavy metals and particles (2019-2023)

Improvements in the Finnish Inventory due to the Inventory Review Processes

EU Technical Review under the NECD in 2021

The improvements made to the 2022 inventory submission in response to the EU Technical Review under the NECD in 2021 are presented in Table 8.2.

Table 8.2 Improvements in response to the 2021 EU Technical Review under the NECD (Final Review Report), actions made in the column to the right

NOTE1 – Responses to NECD Projections Review are provided under the Projections Chapter below.
NOTE2 – Responses to earlier NECD and CLRTAP Reviews can be found in the earlier versions of the IIR.

Review Report Table 4: All findings for CO, PM10 and Black Carbon from the 2021 NECD inventory review 2021

Initial recommendation, year, number of years	KC	Recommendation	RE TC UPTC	Response
FI-1A3biv-2021-0001	No	<p>For 1A3biv Road Transport: Mopeds & Motorcycles, liquid fuels for PM10 and BC in 2017-2018, the TERT noted that, with reference to NFR tables and IIR (chapter transport) pages 14-25, there may be an over-estimate of emissions. This over-estimate does not have an impact on total emissions that are above the threshold of significance. The TERT notes that this issue may be caused by the high increase of PM10 (+21%) and BC (+111%) IEF between 2017 and 2018, whilst fuel consumption decreases by 3%. These trends are not in line with 2000-2017 and 2018-2019 decreasing trends. In response to a question raised during the review, Finland explained that for the 2021 submission the calculation of vehicle category L6e-B under NFR 1A3Biv was thoroughly renewed and recalculated as described in the 2021 IIR under Source specific recalculations. However, in the category L6e-B the recalculation of some compounds for the years 2003-2017 failed. These are PM10 and BC, but failure also concerned PM2.5 and some POPs. Finland provided revised estimates for years 2003-2017 and stated that it will be included in the 2022 submission. The TERT agreed with the revised estimates provided by Finland.</p> <p>The TERT recommends that Finland includes the revised estimates in its 2022 submission.</p>	RE	The revised estimates are included in the 2022 submission.
FI-2D3c-2021-0001 Yes	Yes	<p>For 2D3c Asphalt Roofing and CO, PM10 and BC for 1990-2019, the TERT noted, that the notation key 'NA' (not applicable) is used, whilst a Tier 1 method is available in the 2019 EMEP/EEA Guidebook. In response to a question raised during the review, Finland explained that these emissions are negligible, because abatement systems are installed and 2019 EMEP/EEA Guidebook not always accurately separates process and fuel based emissions. The TERT noted that the issue is below the threshold of significance for a technical correction.</p> <p>The TERT recommends Finland to either estimate emissions or use the Notation Key `NE` (not estimated) and provide an explanation in the IIR, why these emissions are negligible.</p>	No	The explanation was included already in the 2021 IIR under the title of "Particles". To this 2022 IIR the same explanation remains with the title changed into "Particles and CO" on p. 95 of IIR Part 3 IPPU
FI-3B4giv-2021-0001	No	<p>For category 3B Manure management for PM10 and PM2.5 emissions, the TERT noted that that there is a lack of transparency regarding source-specific recalculations in 3B4gi, 3B4gii, 3B4giii, 3B4giv, 3B4h, 3B1b. In response to a question raised during the review, Finland explained that the differences were due to corrections to animal numbers in the 2021 submission. This does not relate to an over- or under-estimate of emissions.</p> <p>The TERT recommends that Finland provides all relevant information regarding recalculations in the next submission.</p>	No	Information on recalculations are included under Chapter "Source-specific recalculations including changes made in response to the review process" in IIR Part 4 Agriculture

Review Report Table 6: All recommendations including those additionally made during the 2021 NECD inventory review and those not implemented from previous reviews, for NOX, NMVOC, SO2, NH3 and PM2.5

Observation	KC	Recommendation	RE, TC or UPTC	Response
FI-3B3-2020-0001	Yes	<p>Assessment of the implementation of the initial recommendation</p> <p>For category 3B3 Manure management – Swine for NH3 emissions, the TERT noted that that Finland uses an average EF for all pig categories (per manure type). This was raised during the 2020 NECD inventory review. In response to a question raised during the review, Finland explained that it is planning to update the agricultural emission calculation system so that it will be possible to use animal specific volatilization coefficients in all animal groups (not only for pigs).</p> <p>The TERT recommends that Finland use animal specific volatilization coefficients in all animal groups for the next submission and provide relevant information in the IIR.</p>	No	The recommendation has been implemented as described in Chapter “Source-specific recalculations including changes made in response to the review process” in IIR Part 4 Agriculture
FI-3Da2c-2020-0001	No	<p>Assessment of the implementation of the initial recommendation</p> <p>For category 3Da2c Other Organic Fertilisers Applied to Soils (including compost) for NOX and NH3 emissions, the TERT noted that there is a lack of transparency regarding the use of the notation key 'NO' while some activity data are provided in the NFR 5B1 Biological treatment of waste – Composting. This was raised during the 2020 NECD inventory review. In response to a question raised during the review, Finland explained that Other organic fertilisers applied to fields, such as composted household waste and industrial waste, were not considered as the amounts applied to fields are considered small. According to the Finnish GHG emission inventory, total nitrogen in other organic fertilizers accounts for approximately 0.3% of the total nitrogen applied in inorganic fertilizers, livestock manure and sewage sludge together. Other organic fertilisers was mainly used in landscaping, not on fields. The TERT noted that the issue is below the threshold of significance for a technical correction.</p> <p>The TERT recommends that Finland either estimates emissions or includes the explanation provided during the review in its next IIR and changes the notation key to 'NE', as this activity is occurring to some extent in Finland.</p>	No	Ammonia and NOx emissions from other organic fertilisers applied to soils were included in the inventory as explained on p. 62 of IIR Part 4 Agriculture.
FI-5B2-2019-0001	No	<p>Assessment of the implementation of the initial recommendation</p> <p>For 5B2 Biological Treatment of Waste - Anaerobic Digestion At Biogas Facilities the TERT noted that NH3 emissions have been reported 'not estimated'. This was raised during the 2019 and 2020 NECD inventory review. The TERT noted that the issue is below the threshold of significance for a technical correction. The 2021 review noted that the IIR states that the issue has been included in the list of improvements and that the recommendation will be addressed in the 2022 submission.</p> <p>The TERT reiterates the recommendation that Finland estimates and reports emissions in the next submission.</p>	No	The emissions are included in the inventory as explained on p. 12-15 of IIR Part 5 Waste

Review Report Table 9: All findings including those additionally made during the 2021 NECD inventory review and those not implemented from previous reviews of LPS data9 Review year of initial data

Initial recommendation (number of years)	Recommendation	Priority recommendation	Response
FI-LPS-GEN-2020-0002	<p>Assessment of the implementation of the initial recommendation</p> <p>The TERT noted that for the year 2019, emissions are reported for 6 facilities in the Industrial Reporting database (v.4) which could not be found in the LPS submission using the EPRTR IDs provided. In response to a question raised during the review, Finland explained that for 4 LPSs, there was an error and these should have been included in LPS reporting, and for the other 2LPSs emission values were below the threshold values (when disaggregating to stack level and excluding some activities). The TERT notes that the 'Guidelines for reporting emission data under the Convention on Long-Range Transboundary Air Pollution' (2015) states that ' "Large point sources" (LPS) are defined as facilities whose combined emissions, within the limited identifiable area of the site premises, exceed the pollutant emission thresholds...' (section III, point 9). The TERT confirms that there is no further deadline for submissions (other than 01-May-2025), but for users of the data (e.g. modelers, scientists) it would be of value to have access to the corrected data sooner.</p> <p>In this context, the TERT recommends that Finland ensures that this issue does not recur in the LPS data submission in 2025 and encourages Finland to provide a corrected version of the current dataset as soon as possible.</p>	Yes	The corrected data will be included in the submission 01-May-2022.
FI-LPS-B-2-2020-0003	<p>Assessment of the implementation of the initial recommendation</p> <p>The TERT noted that for the year 2019, emissions reported for the LPS with the corresponding E-PRTR National ID of 1913 (SSAB Europe Oy (ent. RUUKKI METALS OY), Raahe) are different to those reported for that facility for the same year in the Industrial Reporting database (v.4). In response to a question raised during the review, Finland explained that the 2021 submission of this LPS is likely (partly) incorrect. The TERT confirms that there is no further deadline for submissions (other than 01-May-2025), but for users of the data (e.g. modelers, scientists) it would be of value to have access to the corrected data sooner.</p> <p>In this context, the TERT recommends that Finland ensures that this issue does not recur in the LPS data submission in 2025 and encourages Finland to provide a corrected version of the current dataset as soon as possible.</p>	Yes	The corrected data will be included in the submission 01-May-2022.

Review Report Table 11: All findings including those additionally made during the 2021 NECD inventory review and those not implemented from previous reviews for gridded data

Observation	Recommendation	Priority recommendation	Response
FI-GRID-GEN-2020-0003	<p>Assessment of the implementation of the initial recommendation For the gridded data submitted for the year 2019 the TERT noted there are a number of locations (grid cells) where gridded and LPS data are inconsistent. The TERT had compared gridded emissions for each grid cell with LPS emissions (allocated to the respective grid cell), where several inconsistencies were found, i.e. cases where LPS emissions exceeded gridded emissions. This was previously raised during the 2020 NECD inventory review and is therefore flagged as a priority recommendation. In response to a question raised during the review Finland explained that there was a mistake in the gridded data file (see also observation FI-GRID-A-2020-0001) and a resubmission was provided by Finland. Due to time constraints during the review, the TERT was unable to check the consistency of LPS and gridded based on the resubmitted data.</p> <p>The TERT recommends that for the next submission, Finland ensures the gridded data are consistent with reported LPS emissions.</p>	Yes	The coordinates of facilities under LPS and gridded emissions has been synchronized in the submission of 2021.
FI-GRID-A-2020-0001	<p>Assessment of the implementation of the initial recommendation The TERT noted, with reference to gridded emissions for 2019, an issue in the gridded emissions which caused strong discrepancies with the national inventory totals for all pollutants. In response to a question raised during the review, Finland provided a resubmission which resolved nearly all discrepancies. However, since also during the 2020 NECD inventory review multiple submissions provided by Finland prior to and during the review contained serious problems regarding the spatial variability across all pollutants, the TERT notes a pattern of mistakes and improper quality control of the data prior to submission. Therefore, the issue will be flagged as a priority recommendation.</p> <p>The TERT recommends that for the next submission, Finland should carefully check the gridded data (the Annex V template) prior to the submission in order to prevent these types of error.</p>	Yes	An added layer of quality control will be added in the 2022 submission of gridded data in order to ensure data integrity during submission process.

8.2 PROJECTIONS

Changes in chapter	
Update of text March 2022	KS, MS
Update of projections	Every 1-3 years, since 2020 every 2 years

8.2.1 Projections for 2020, 2025 and 2030

With existing measures (WM) projections


The activity pathways in Finland's Energy and Climate Strategy (Huttunen, 2017¹) are used as the basis for most of the projections. The Strategy had two scenarios: With Existing Measures and With Additional Measures. When making the National Air Pollution Control Programme 2030 (Ministry of the Environment, 2019²), the two scenarios were combined to best represent the contemporary status of national legislation and agreed measures. Emissions for this one combined scenario, called Baseline, were then estimated. It was found that the Baseline scenario is already expected to meet the reduction targets for Finland. This conclusion was further supported by a sensitivity analysis for key sectors. Due to this, no additional scenarios were made (although the Programme did suggest additional measures). The Baseline scenario of the Programme is called WM scenario in this document. Some tweaks have been made to the projections with the emergence of new information (e.g. emission factors for some wood burning appliances), but a more thorough update for most sectors will follow in 2023 after a new Energy and Climate Strategy has been completed. Projections for the transport sector presented here are already in line with the upcoming Energy and Climate Strategy.

Projections for 2020, 2025, 2030, 2040 and 2050

Emission projections for 2020, 2025 and 2030 submitted in 2022 cover nitrogen oxides, sulphur oxides, non-methane volatile organic compounds, ammonia, fine particles (PM2.5) and black carbon.

For agriculture and transport sectors, emission scenarios are available up to 2050. For IPPU, fugitive emissions and waste sector, expert estimates were made also for the years 2040 and 2050 for the 2020 submission, however, these will be further developed in the coming years. The current projected emission values are presented in Table 8.3.

Table 8.3. Projected national total emissions for 2015, 2020 and 2030 as reported on 15 March 2022



¹ Huttunen R. (Ed.) 2017. Government Report on the National Energy and Climate Strategy for 2030. Publications of the Ministry of Economic Affairs and Employment, 2017-02-08.

² Ministry of the Environment. 2019. National Air Pollution Control Programme 2030. Publications of the Ministry of Environment 2019:7

8.2.2. Projections for Energy

Changes in chapter	
Update of text March 2022	MS, KS
Update of projections	Every 1-3 years, since 2020 every 2 years

Projection for energy are based on the development of emissions from 2020 to 2030 in the National Air Pollution Control Programme 2030. In the Programme, the FRES model (Karvosenoja 2008³) was used to calculate emission projections for combustion plants, industrial processes and residential combustion. It covers the emissions of sulfur dioxide (SO₂), nitrogen oxides (NO_x), ammonia (NH₃), non-methane volatile organic compounds (NMVOCs) and primary particulate matter (TSP, PM₁₀, PM_{2.5}, PM₁ and PM_{0.1}). Primary PM includes the fractionation to main chemical species (black and organic carbon, sulfate, main heavy metals and mineral matter).

FRES is a scenario model, where a new reference year is added every five years (2010, 2015 etc.) and target years are selected according to specific needs and available activity data. The intermediate years are presented linearly. Since 2015 was the most recent historic year when making the National Air Pollution Control Programme 2030, it was used as the reference year at the time. However, here the reported 2020 emissions are used as the reference, and projection made with the FRES model have been used to estimate the relative change in emissions from 2020 to 2030. This has been done by applying the relative change of the projections between 2020-2030 to the reported emissions of 2020.

Model parameters

The emissions are calculated from the parameters of activity levels, emission factors and emission control technology removal efficiencies and utilization rates (Figure 8.2). The energy consumption and industrial production scenarios produced in the national Energy and Climate Strategy are used as input to the model (Annex IV B-WM. The reference year in cells B10-B40 is 2014, since that was the reference year in the Strategy). In the FRES model the activity unit for combustion processes is annual primary energy use (e.g. PJ a-1). Emission sources are treated as point sources (~450 combustion plants and ~130 process industry plants) or area sources (residential combustion and small or less active combustion plants).

For some combustion plants the emissions and fuel use data are reported in the national YLVA database so that plant-specific emission factors can be calculated. If the data is not available, emission factors will be implemented based on legislation (current or upcoming, depending on the year). The statutes affecting the emission limits of combustion plants are:

- The Industrial Emissions Directive, and the BAT conclusions concerning energy production and different industrial sectors, Medium Combustion Plant Directive
- Environmental Protection Act (527/2014)
- Government Decree on Limiting Emissions from Large Combustion Plants (936/2014)
- Government Decree on Environmental Protection Requirements for Medium-sized Energy Production Units (1065/2017)
- Government Decree on Waste Incineration (151/2013)

Based on these a table has been compiled, where emission factors are given for each fuel type and plant size category, taking into account the age of the plant (available at request).

Small-scale residential combustion is the biggest source of PM and NMVOC emissions in Finland. The emission calculation scheme for residential (wood) combustion is described in Savolahti et al. (2016)⁴. It

³ Karvosenoja N. 2008 Emission scenario model for regional air pollution. *Morographs Boreal Environ. Res.* 32. 2008.

⁴ Savolahti M., Karvosenoja N., Tissari J., Kupiainen K., Sippula O. & Jokiniemi J. 2016. Black carbon and fine particle emissions in Finnish residential wood combustion: Emission projections, reduction measures and the impact of combustion practices. *Atmospheric Environment* 140 (2016) 495-505. <https://doi.org/10.1016/j.atmosenv.2016.06.023>

includes 5 categories for small-scale central heating boilers and 9 categories for stoves or fireplaces. All of them have separate emission factors based mostly on national measurements, and annual activities based on questionnaires. We have also tried to take into account the suboptimal combustion practices of some stove users. Based on emission factors from literature and info from chimney sweeps, we have used a coefficient for “poor combustion”, resulting in an increase of the average emission factors. Future emissions in the projection are determined by activity changes, natural development of the appliance stock and the Ecodesign directive (2015/1195 and 2015/1189 for residential combustion). The prevalence of wood combustion has been increasing during the last decades, and this trend is expected to continue up to 2030, although improvements in energy efficiency are expected to reduce the overall heating demand. Ecodesign will not have a major impact until 2030, since it mainly targets appliances with a very long service life (e.g. ~35 years for masonry heaters) and does not cover sauna stoves, which are a major polluter in Finland. However, the natural development of the appliance stock towards cleaner stoves and boilers is expected to notably decrease the average emission factors of the sector by 2030, and thus decrease the emissions.

For ammonia emissions in the energy sector, the projections are expert estimates based on knowledge of fuel use at plants.




Figure 8.2. Structure of the FRES model.

8.2.3 Projections for Transport

Changes in chapter	
Update of text March 2022	AL, JoMa, KS
Update of projections	Every 1-3 years, since 2020 every 2 years

Projections for the transport sector are presented in Figure 8.3.




Figure 8.3 Projections for the transport sector

Projections for road and rail transport as well as aviation and working machines are produced using VTT's LIPASTO system, which has a time series for the years 1980-2050.

Road transport

In road transport, the projections are based on the authorities' (The Finnish Transport Infrastructure Agency) forecast of vehicle kilometrage until 2050⁵. The vehicle kilometrage forecast is affected by several factors: most importantly, forecasted development of economy, population growth and cost of driving. A major change in methodology is the recently increased passenger kilometrage: the relatively inexpensive driving of battery electric vehicles is assumed to increase passenger vehicle kilometrage. In the ALIISA model, average annual kilometrages for vehicles are adjusted so that the national kilometrage forecast is achieved. Thus, the composition of fleet (vehicle types and motive powers) is linked to the projected kilometrage development. Sales forecasts cover both new vehicles and used vehicle imports, which in Finland is a significant amount. Scrappage rate is based on an estimate based on actual changes in the fleet. The model considers the penetration of the Euro classes based on statistical data and the fuel efficiency development of the vehicles according to the CO₂ emission regulation by EU⁶. Fuel efficiency of new vehicles is assumed to be improved mainly due to increased sales of plug-in-hybrid and battery electric vehicles.

The modelled fleet is divided into 40 different sub-types: 5 main types, passenger cars (with and without catalytic converter), vans, buses, trucks without a trailer and trucks with a trailer. These, in turn, are divided into seven propulsion groups: gasoline, diesel, E85, ED95, gas, electricity and hydrogen. Each of these have their own forecasts of shares of annual sales and imports, and fuel efficiency development. In addition, regarding consumed fuels, the proportions of fossil and renewable fuel components are considered. The use of biofuels will increase from 20 % (by energy content, of annual liquid fuel sales) up to 30 % during the period 2020-2029 due to the biofuel obligation in Finland⁷. Biogas is included in the obligation since 2022, but not included in the forecast.

The baseline⁸ forecast is that new passenger car sales will be on average 4.7% annually in 2016-2020, 4.4% on average in 2021-2030 and 4.7% on average in 2031-2050. The corresponding figures for vans are: 4.8% 4.8% 4.8% and for trucks: 4.0% 4.0% 3.9%. A special feature in Finland are the very heavy trucks (gigatrucks), which reduce the number of trucks with trailer. The increase in the number and kilometrage of motorcycles and mopeds is expected to stabilise in the coming years. The vehicle fleet development is based on an expert estimate on the current sales trends of available powertrains and the availability of new vehicles in the European market. The development is dominated in the forecast by battery electric vehicles, which replace conventional gasoline and diesel vehicles in all vehicle categories except in trucks, for which CNG/LNG-powered vehicles replace conventional vehicles. The demand for electric vehicles is high, but a short supply limits the expected pace of the fleet turnover.

Ammonia

In transport, most NH₃ emissions originate from passenger cars equipped with catalytic converters. Improvements in technology have substantially reduced NH₃ emissions from passenger cars after 2005 and the development is continuing in the projections. On the other hand, the introduction of the urea additive in heavy vehicles since Euro V significantly increases their ammonia emissions. However, since heavy-duty NH₃ emissions are one-tenth of NH₃ emissions from passenger cars, emissions from passenger cars dominate and overall emissions are decreasing in the projections.

⁵ Finnish Transport and Communications Agency (Traficom). 2021. Tieliikenteen valtakunnallinen ennuste (in Finnish). Available at: https://api.hankeikkuna.fi/asiakirjat/d99a3ae3-b7f9-49df-afd2-c8f2efd3dc1d/5d4db40f-4cf4-4ad5-9e42-99031175f27e/MUISTIO_20210920060527.pdf

⁶ Regulation (EU) 2019/631 of the European Parliament and of the Council of 17 April 2019 setting CO₂ emission performance standards for new passenger cars and for new light commercial vehicles

⁷ Finnish Act on biofuels 13.4.2007/446. Available at: <https://www.finlex.fi/fi/laki/ajantasa/2007/20070446>

⁸ Ministry of Transport and Communications & Technical Research Centre of Finland VTT Ltd. 2021. Baseline emissions for transport until 2050. Result excel-workbook available at: https://api.hankeikkuna.fi/asiakirjat/d99a3ae3-b7f9-49df-afd2-c8f2efd3dc1d/1a483d38-ee85-418e-9086-fd5ef6ea40d0/MUISTIO_20210920060401.docx

PM 2.5 and BC

Car exhaust emissions of particulate matter (PM_{2.5}) and black carbon (BC) have fallen to less than a quarter since the year 2000 and the development is continuing in the projections. However, emissions from tyre and brake wear and road abrasion are increasing. According to the forecast, the vehicle kilometrage for passenger cars especially, increases from 2020 until 2050. One key reason for this is relatively inexpensive driving of battery electric vehicles.

NMVOG

NMVOG emissions are decreasing due to the renewal of vehicle fleet. A significant share of these emissions is caused by mopeds and motorcycles. The increase in the number and kilometrage of motorcycles and mopeds is expected to stabilise in the coming years and the vehicle fleet renewal towards greater share of newer Euro-classes is continuing in the projections. Currently there is no detailed information on the development of NMVOG emissions in the coming years.

Railways

The forecast for rail transport is based on the expert estimation of the development of diesel train transport volume. The proportion of diesel trains has decreased significantly in recent years and they are mainly used in non-electrified, smaller rail sections. The diesel train transport has stabilised, and no major changes are expected.

National navigation

In maritime transport, the calculation of the MEERI model is mainly based on the number of port calls at Finnish ports. The forecasts therefore focus on experts' (The Finnish Transport Infrastructure Agency's) estimation of the development of port calls. Emission factor forecasts are based on expert estimation on the development of different Tier emission levels in ships and the use of different fuels (HFO, HFO with scrubbers, MDO/MGO, diesel, LPG). For icebreakers, where emissions are dependent on highly changing yearly ice conditions, forecasts are based on a 10-year average.

In work vessels, ferry boats, fishing vessels and leisure boats the situation has been stabilised and no changes are expected.

Working machines

For working machines, the forecasts are based on a recent analysis of alternative powertrains and their market penetration in Finland⁹. Evolution of fleet (50 different machine types) is affected by, the penetration of emission standards (Stage levels) for ICE-powertrains and the market penetration of electric work machines in Finland. Mechanisation of the work has reached its maximum and the sales of new machines is mainly replacing scrapped machines and the number of most of the machine types is stabilised. Increased efficiency of machines, work and emission restrictions and increasing electrification of machines will reduce emissions in projections.

Aviation

There is currently no model to calculate aviation emissions, which are minor and are not considered to impact the projected trends.

⁹ Markkanen, J., & Lauhkonen, A. (2021). Työkoneiden päästöjen perusennuste ja sähköistymisen vaikutus päästöihin (in Finnish). VTT Technical Research Centre of Finland. VTT Asiakasraportti No. VTT-CR-00245-21. Available at: <https://cris.vtt.fi/en/publications/ty%C3%B6koneiden-p%C3%A4st%C3%B6jen-perusennuste-ja-s%C3%A4hk%C3%B6istymisen-vaikutus-p%C3%A4st%C3%B6ihin>

8.2.4. Projections for IPPU and Waste

Changes in chapter	
Update of text March 2022	KS, MS
Update of projections	Every 1-3 years, since 2020 every 2 years

Projections for the IPPU sector were updated based on best available information depending on the activity sector as described below in Table 8.4.

For those emissions indicated below to be based on the FRES model, the reference year, we have used reported emissions (average of preceding years) of the industrial facilities as found in the national YLVA database. The projections are a combination of assumed activity changes and developments in cleaner technology. A specific inquiry of the future development prospects was conducted to main industrial sectors as part of the National Air Pollution Control Programme 2030

Table 8.4. Methods and assumptions used in IPPU and Waste sector projections

Sector		Method
2A1	Cement production	<p>NMVOC emissions occur from 2A1 and 3A3 but are minor and are considered to be already included in the uncertainty of the air pollutant emissions from the major IPPU sectors.</p> <p>NOx from 2A2 and 2A3 are included in the energy sector projections as they also are in the inventory of historic years.</p> <p>All NOx, SOx, PM2.5 and BC projections are estimated in the FRES model based on average of preceding years and are a combination of assumed activity changes and developments in cleaner technology. A specific inquiry of the future development prospects was conducted to main industrial sectors as part of the National Air Pollution Control Programme 2030.</p>
2A2	Lime production	
2A3	Glass production	
2A5a	Quarrying and mining of minerals other than coal	
2A5b	Construction and demolition	
2A5c	Storage, handling and transport of mineral products	
2B2	Nitric acid production	
2B6	Titanium dioxide production	
2B10a	Chemical industry: Other	
2B10b	Storage, handling and transport of chemical products	
2C1	Iron and steel production	
2C2	Ferroalloys production	
2C3	Aluminium production	
2C6	Zinc production	
2C7a	Copper production	
2C7b	Nickel production	
2C7c	Other metal production	

2C7d	Storage, handling and transport of metal products	
2D3a	Domestic solvent use including fungicides	NMVOC projections are calculated using capita forecasts as a surrogate
2D3b	Road paving with asphalt	NMVOC projections are based on the running average of last three years. No forecast is available for the development of the sector.
2D3c	Asphalt roofing	
2D3d	Coating applications	
2D3e	Degreasing	
2D3f	Dry cleaning	
2D3g	Chemical products	
2D3h	Printing	
2D3i	Other solvent use	
2G	Other product use	
2H1	Pulp and paper industry	SOx, PM2.5 and BC emissions projections are estimated in the FRES model based on average of preceding years and are a combination of assumed activity changes and developments in cleaner technology. A specific inquiry of the future development prospects was conducted to main industrial sectors as part of the National Air Pollution Control Programme 2030 NMVOC and NH3 emission projections are based on the running average of last three years.
2H2	Food and beverages industry	NH3 projection is based on a running three year average of in relation to capita forecasts for the projected years.
2I	Wood processing	NMVOC emission projection is based on the running average of last three years. No forecast is available for the development of the sector.
2L	Other production, consumption, storage, transportation or handling of bulk products	IPPU sector emissions are minor and are thus considered to be included in the uncertainty of the air pollutant emissions from the other IPPU sectors.
5A	Biological treatment of waste - Solid waste disposal on land	A minor and constantly decreasing source of NMVOC and particle emissions due to constantly increasing incineration of waste with energy recovery. The emissions are minor and are thus considered to be included in the uncertainty of the air pollutant emissions from the other IPPU sectors.
5B1	Biological treatment of waste - Composting	A minor source of air pollutants, estimated to be constant
5B2	Biological treatment of waste - Anaerobic digestion at biogas facilities	The forecast for NH3 is estimated based on principles explained below for the agriculture sector projections.
5C1bv	Cremation	Ox , SOx and NMVOC projections from 2G are included in the energy sector projections as they also are in the inventory of historic years.

		PM2.5 and BC emissions are minor and are thus considered to be included in the uncertainty of the emissions from the other IPPU sectors.
5D1	Domestic wastewater handling	NMVOC emissions are minor and are considered to be included in the uncertainty of the emissions from the other IPPU sectors. NH3 projection is based on a running three year average of in relation to capita forecasts for the projected years.
5D2	Industrial wastewater handling	NMVOC emissions are minor and are considered to be included in the uncertainty of the emissions from the other IPPU sectors.
5E	Other waste	PM2.5 and BC emissions are minor and are considered to be included in the uncertainty of the emissions from the other IPPU sectors.

Population forecasts, GDP, GDP/capita, GCP forecasts and GDP growth forecasts used in the preparation of the IPPU and Waste sector projections are presented in Table 8.5. The figures in orange italics are calculated from the latest data available (2020) with GDP growth forecast in relation to population forecasts.

In the projections the structure changes in the Finnish economy 1075-2020 (Figure 8.4) are taken into account, especially the latest years decreasing trends due to covid and the Ukrainian war, which is likely to impact negatively to IPPU activities in Finland. In addition, for decades, the emission levels in Finland have decreased even when the activity levels have increased, due to implementation of BAT technologies.

Table 8.5 Population forecasts, GDP, GDP/capita, GCP forecasts and GDP growth forecasts

Year	Population (mill.) (Statistics Finland)	GDP (Statistics Finland)	GDP/capita (Statistics Finland)	GDP growth forecast ¹⁾ ¹⁰
1980	4.8	33657	7030	
1990	5.0	90959	18197	
2000	5.2	136442	26334	
2007	5.3	187072	35293	
2010	5.4	188143	35002	
2013	5.5	204321	37481	
2016	5.5	217518	39525	
2019	5.5	239852	43410	
2020	5.5	236032	42653	
2021	5.5	<i>244293</i>	<i>44040</i>	3.5
2022	5.6	<i>254065</i>	<i>45736</i>	4.0
2023	5.6	<i>257368</i>	<i>46268</i>	1.3
2024	5.6	<i>260456</i>	<i>46764</i>	1.2
2025	5.6	<i>263582</i>	<i>47269</i>	1.2
2030	5.6	<i>266745</i>	<i>47643</i>	1.2
2050	5.5	<i>269419</i>	<i>48779</i>	0.5
2060	5.4	<i>270766</i>	<i>49707</i>	0.5

¹⁰ Pellervo ja Euro ja Talous <https://www.eurojatalous.fi/fi/2021/5/suomen-uuden-pitkan-aikavalin-ennusteen-mukaan-kasvunayntaa-entista-vaisummalta/>




Figure 8.4. Structural changes in Industry (% of brutto value addition): Services (blue line), Industry and construction (orange line), Agriculture, Forestry and Fishing industry (red line)

8.2.5 Projections for Agriculture

Changes in chapter	
Update of text March 2022	JG, KS
Update of projections	Every 1-3 years, since 2020 every 2 years

Projections for agriculture are based on the national Agriculture sector calculation model available up to 2050. The animal numbers, development of nitrogen excretion and mineral fertilizers and land use areas are forecasts by LUKE based on the Dynamic Regional Sector Model of Finnish Agriculture, Dremfia, except for fur animals and reindeer, for which the numbers are estimated from existing statistics assuming that there will not be major changes in the coming years. (Lehtonen, 2021¹¹).

There are some differences in projected emissions for agriculture compared to the ones presented in the previous reporting. The reasons for these differences are:

- Changes in animal number and inorganic fertilizer use estimations: for cattle, sheep and goats, animal statistics for 2020 were already available when the projections were calculated and they were used. The numbers of goats and suckler cows are higher compared to the estimated ones, whereas the numbers of all other cattle and sheep are lower. Additionally, the animal number estimations for the other animal groups for 2020 and for all animal groups for the years 2025, 2030, 2040 and 2050 were updated based on the data provided by the Natural Resources Institute Finland (LUKE; Lehtonen 2021). Furthermore, the annual inorganic nitrogen fertilizer consumption estimations were updated by Lehtonen (2021). Animal numbers and the use of mineral nitrogen fertilizers are the same as are used in the latest national ghg-emission WEM-scenarios for agriculture.
- Changes in estimated nitrogen excretion rates: nitrogen excretion rates of bulls, dairy cows, heifers, suckler cows and weaned pigs were re-estimated based on the recent development. For all above mentioned animal groups the N excretion rate was estimated to be higher in the projected years than was estimated earlier.
- Changes in manure management:
 - o the share of injected slurry was lowered (70% -> 65%) for 2020, based on the most recent knowledge,
 - o shares of slurry and urine incorporation methods (for slurry and urine spread on stubble) were corrected for 2025, 2040 and 2050,

¹¹ Lehtonen, H. 2021. Personal communication 12.2.2021. Natural Resources Institute Finland (LUKE)

- for pig slurry it was estimated that cooling of slurry channels and increased slurry removal frequency will be more common in the future, due to the implementation of IRPP BAT conclusions in pig production.
- A ban on field burning of agricultural residues from the beginning of 2021, resulting zero emissions for the year 2025 and beyond.

Activity data for agricultural emission projections

- *Animal numbers* and inorganic nitrogen use data used in the emission projections are shown in the tables 8.6-8.19.
- Estimated evolution of *manure management practices* is described in the tables below. Efficiency of the emission abatement measures (as well as the unabated emission factors used) can be found from the Table 5.11 of the Agriculture-chapter of the IIR.
- *Nitrogen excretion rate estimations* are based on the recent development and are shown in in the tables below.

The development of animal numbers, the use of mineral fertilizers and land use areas are projected up to 2050 in Natural Resources Institute Finland (Luke) based on the Dynamic Regional Sector Model of Finnish Agriculture, Dremfia. This model is based on agricultural economics principles, is well validated in terms of production and land use 1995-2020, and has been frequently used in analysing agricultural and agri-environmental policies (Lehtonen & Niemi 2018, Lehtonen & Niskanen 2016, Lehtonen & Rankinen 2015). However, the development of the numbers of fur animals, lambs, goats and reindeers, which have a small economic role and a small contribution to e.g. greenhouse gases and ammonia, are estimated from existing statistics assuming that there will not be major changes in the coming years. These assumptions and principles have been used when projecting development of agriculture and greenhouse gases in Finland in recent years (see e.g. Aakkula et al. 2019, Koljonen et al. 2020, Lehtonen & Rämö 2020).

The DREMFA sector model has produced several times projections of animal numbers which fit very well the post-development 1995-2020. In fact, the animal number projections in the references above, shows decreasing numbers of bovine animals and pigs, but non-decreasing numbers of poultry animals for years after 2020. This is because of the increasing milk yield of dairy cows which imply reduced numbers of other bovine animals, and because the number of suckler cows is very unlikely to increase due to high dependence on farm subsidies in suckler cow production and non-increasing coupled payments for bovine animal production. Pig numbers have been decreasing since ten years due to gradually decreasing domestic demand of pig meat and reduced profitability of production while consumers have substituted poultry meat for pork and thus the number of poultry animals have been in the increase. There is a reason to assume, based on healthiness and climate awareness of consumers, that red meat production (beef, pork) in Finland will continue on gradually decreasing trend and thus the numbers of bovine animals and pigs will remain below the current levels in the future.

Note on 3F, Field burning and other agriculture projections: *Field burning is forbidden starting from 1.1.2021*

Table 8.6. Animal numbers and use of inorganic nitrogen fertilisers in 2015-2020 and as prediction for 2025-2050

	Cattle					Fur Animals			Horses and ponies		Pigs			
	Bull >1 yr	Calf <1 yr	Dairy cow	Heifer >1 yr	Suckler cow	Fox and racoon	Mink and fitch	Reindeer	Horse	Pony	Boar (50- kg)	Fattening pig (50-kg)	Sow (with piglets)	Weaned pig (20-50 kg)
Finland 2015	109 348	306 899	285 147	154 610	58 713	2 084 444	1 026 774	191 100	63 800	10 400	2 100	501 100	345 300	115 800
Finland 2016	107 786	309 650	282 443	150 158	58 984	2 202 072	1 027 352	191 473	63 800	10 400	1 600	489 200	342 300	108 100
Finland 2017	110 767	297 334	274 954	150 269	59 854	2 027 757	933 504	193 142	64 000	10 400	1 400	446 900	312 400	99 100
Finland 2018	105 770	298 882	271 429	146 100	60 096	2 017 309	842 509	184 958	64 000	10 400	1 400	425 000	273 300	95 200
Finland 2019	104 485	288 066	262 292	142 430	60 349	1 889 369	679 061	188 190	63 900	10 400	1 300	438 600	272 400	92 200
Finland 2020	98 187	290 199	259 579	136 445	61 975	1 295 961	500 498	188 190	64 000	10 000	1 100	450 600	299 700	88 700
Finland 2025	93 299	275 717	237 891	129 688	67 659	2 000 000	1 000 000	190 000	70 507	11 493	1 299	394 318	253 570	88 327
Finland 2030	89 042	263 135	230 504	123 770	61 102	2 000 000	1 000 000	190 000	70 507	11 493	1 266	384 181	247 051	86 057
Finland 2040	83 406	246 480	207 447	115 936	65 702	2 000 000	1 000 000	190 000	70 507	11 493	1 230	373 448	240 149	83 652
Finland 2050	79 352	234 501	195 941	110 301	63 933	2 000 000	1 000 000	190 000	70 507	11 493	1 212	368 006	236 649	82 433

	Poultry							Goat and sheep		Inorg. N
	Broiler breeder hen	Broiler	Laying hen pullet	Cockerel	Laying hen	Other poultry	Turkey	Goat	Sheep	Use of inorganic N-fert. as N (tonnes)
Finland 2015	548 175	7 827 340	662 228	25 500	3 594 537	23 246	245 906	4 546	155 238	143 479
Finland 2016	523 216	8 271 563	747 625	26 271	3 598 941	16 930	260 311	4 799	156 496	138 128
Finland 2017	472 976	8 046 698	508 874	22 335	3 745 944	47 213	291 579	5 278	155 926	138 948
Finland 2018	424 296	8 780 903	607 630	17 426	3 984 824	26 135	299 093	5 437	154 999	138 385
Finland 2019	394 715	9 111 743	647 260	16 405	3 900 385	26 758	262 646	5 925	144 876	146 798
Finland 2020	396 097	8 507 327	566 326	17 958	3 811 547	9 639	267 986	6 034	149 171	139 316
Finland 2025	544 667	9 109 724	547 638	22 027	3 591 556	20 000	310 293	6 100	140 200	135 280
Finland 2030	543 312	9 087 055	527 505	21 218	3 459 513	20 000	309 521	6 100	140 200	136 712
Finland 2040	548 664	9 176 581	491 374	19 764	3 222 560	20 002	312 570	6 100	140 200	137 261
Finland 2050	549 097	9 183 815	459 876	18 503	3 016 986	20 004	312 817	6 100	140 200	137 691

Table 8.7 Nitrogen excretion rates (kg N/animal place/year) in 2015-2020 and as prediction for 2025-2050

		Dairy cow	Heifer >1 yr	Calve <1 yr	Suckler cow	Bull >1 yr	Sow	Boar	Fattening pig	Weaned pig	Horse	Pony	Sheep	Goat
Finland	2015	131,910	54,985	40,441	75,669	68,666	31,374	20,562	17,385	9,117	61,913	44,329	7,260	10,700
Finland	2016	130,419	54,585	40,281	75,028	68,714	32,060	20,558	17,275	9,117	61,987	44,515	7,310	10,700
Finland	2017	132,830	55,507	40,906	74,877	69,633	31,952	20,813	17,217	9,131	61,972	44,476	7,430	10,700
Finland	2018	133,374	55,659	41,136	75,303	70,647	32,876	21,025	17,206	9,162	61,898	44,593	7,600	10,700
Finland	2019	136,712	57,802	42,811	77,276	73,012	32,268	20,941	17,170	9,162	61,769	44,681	7,600	10,700
Finland	2020	139,062	58,952	43,705	78,355	74,788	33,794	20,692	17,064	9,177	61,664	44,693	7,600	10,700
Finland	2025	143,000	60,000	44,500	79,000	75,000	32,060	20,558	17,275	9,150	61,987	44,515	7,600	10,700
Finland	2030	150,000	62,000	46,000	80,500	77,000	32,060	20,558	17,275	9,150	61,987	44,515	7,600	10,700
Finland	2040	160,000	66,000	50,000	84,000	82,000	32,060	20,558	17,275	9,150	61,987	44,515	7,600	10,700
Finland	2050	160,000	66,000	50,000	84,000	82,000	32,060	20,558	17,275	9,150	61,987	44,515	7,600	10,700

		Broiler	Broiler hen	Laying hen	Laying hen pullet	Cockerel	Turkey	Other poultry	Fox and racoon	Mink and fitch	Reindeer
Finland	2015	0,478	0,992	0,623	0,388	0,969	1,629	0,636	3,000	1,305	10,700
Finland	2016	0,479	0,992	0,615	0,388	0,969	1,629	0,636	3,000	1,305	10,700
Finland	2017	0,478	0,992	0,601	0,389	0,969	1,663	0,636	3,000	1,305	10,700
Finland	2018	0,478	0,992	0,601	0,389	0,969	1,640	0,636	3,000	1,305	10,700
Finland	2019	0,478	0,992	0,583	0,388	0,969	1,695	0,636	3,000	1,305	10,700
Finland	2020	0,479	0,992	0,593	0,388	0,969	1,700	0,636	3,000	1,305	10,700
Finland	2025	0,477	0,992	0,590	0,388	0,969	1,650	0,636	3,000	1,305	10,700
Finland	2030	0,477	0,992	0,590	0,388	0,969	1,650	0,636	3,000	1,305	10,700
Finland	2040	0,483	0,992	0,590	0,388	0,969	1,650	0,636	3,000	1,305	10,700
Finland	2050	0,483	0,992	0,590	0,388	0,969	1,650	0,636	3,000	1,305	10,700

Manure management data for 2020 and as a prediction for 2025, 2030 and 2040 (in most of the cases, values used for 2050 are the same as are used for 2040).

All values: % of manure unless otherwise stated.

Table 8.8. General information on cattle manure management and grazing in Finland in 2020 and as a prediction for 2025, 2030 and 2040.

	Dairy cows				Suckler cows				Heifers				Bulls				Calves <1 yr			
	2020	2025	2030	2040	2020	2025	2030	2040	2020	2025	2030	2040	2020	2025	2030	2040	2020	2025	2030	2040
<u>Manure management</u>																				
Treated as slurry (%)	78	78	78	78	6	4	1	1	59	60	65	65	59	55	50	40	44	45	50	50
Treated as deep litter (%)	3	3	3	3	35	37	40	40	7	7	5	5	11	15	20	30	15	15	15	15
Treated as solid manure (%), of which	19	19	19	19	59	59	59	59	34	33	30	30	30	30	30	30	41	40	35	35
urine not separated (FYM, %)	23	23	23	23	91	91	91	91	58	58	58	58	88	88	88	88	63	63	63	63
urine separated (%)	77	77	77	77	9	9	9	9	42	42	42	42	12	12	12	12	37	37	37	37
<u>Grazing</u>																				
Grazing period (days)	138	138	138	138	171	171	171	171	134	134	134	134	161	161	161	161	127	127	127	127
Grazed animals (%)	73	73	73	73	92	90	90	90	68	65	65	65	9	9	9	9	31	30	30	30
Animals inside in nights (%)	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100
Time inside at night (h)	12	12	12	12	1	1	1	1	1,5	1,5	1,5	1,5	1	1	1	1	2	2	2	2
Manure excreted on pasture (%)	14	14	14	14	41	40	40	40	23	22	22	22	4	4	4	4	10	10	10	10

Table 8.9. General information on manure management and grazing of sheep, goats, horses, ponies, fur animals and reindeer in Finland in 2020 and as a prediction for 2025, 2030 and 2040.

	Sheep & Goat 2020-2040	Horses & Ponies 2020-2040	Fur animals 2020-2040	Reindeer 2020-2040
<u>Manure management</u>				
Treated as slurry (%)	0	0	0	-
Treated as deep litter (%)	50	13	0	-
Treated as solid manure (%), of which	50	87	100	-
urine not separated (FYM, %)	100	100	100	-
urine separated (%)	0	0	0	-
<u>Grazing</u>				
Grazing period (days)	153	180	-	365
Grazed animals (%)	90	97	-	100
Animals inside in nights (%)	100	100	-	0
Time inside at night (h)	2	6	-	0
Manure excreted on pasture (%)	35	36	-	100

Table 8.10. Percentages (%) of management methods for pig manure in Finland in 2020 and as a prediction for 2025, 2030 and 2040.

	Sows				Fattening pigs 2020-2040	Boars 2020-2040	Weaned pigs 2020-2040
	2020	2025	2030	2040			
Treated as slurry (%)	90	85	80	70	100	90	90
Treated as deep litter (%)	0	5	10	20	0	0	4
Treated as solid manure (%), of which	10	10	10	10	0	10	6
urine not separated (FYM, %)	30	30	30	30	18	30	11
urine separated (%)	70	70	70	70	82	70	89

Table 8.11. Percentages (%) of management methods for poultry manure in Finland in 2020 and as a prediction for 2025, 2030 and 2040.

	Laying hens				Broilers 2020 – 2040	Laying hen pullets				Cockerels 2020-2040	Broiler hens 2020 - 2040	Turkeys 2020 - 2040	Other poultry 2020- 2040
	2020	2025	2030	2040		2020	2025	2030	2040				
Slurry	0	0	0	0	0	0	0	0	0	0	0	0	0
Deep litter	10	25	25	25	100	40	40	40	40	50	100	100	40
Solid manure	90	75	75	75	0	60	60	60	60	50	0	0	60

Table 8.12. Detailed information on slurry management in animal shelters and manure storages in Finland in 2020 and as a prediction for 2025, 2030 and 2040. Unit: percentage of total slurry per animal species.

Abatement measures	Slurry											
	Cattle				Pigs				Poultry (no slurry systems in 2020 or later)			
	2020	2025	2030	2040	2020	2025	2030	2040	2020	2025	2030	2040
<u>Animal shelter (% of manure)</u>												
Improved cleaning of surfaces	15	17	20	25	20	20	20	20	0	0	0	0
Flushing	2	2	2	2	4	4	4	4	0	0	0	0
Increased manure removal frequency	4	4	4	4	10	15	30	50	0	0	0	0
Rapid urine separation	-	-	-	-	-	-	-	-	-	-	-	-
Biological or chemical air scrubbers	0	0	0	0	2	2	2	2	0	0	0	0
Cooling of manure channels	0	0	0	0	15	20	25	50	-	-	-	-
Drying of manure on manure belt	-	-	-	-	-	-	-	-	-	-	-	-
Non-leaking drinking system	-	-	-	-	-	-	-	-	-	-	-	-
<u>Manure storage (% of manure)</u>												
No measures	0	0	0	0	27	23	15	15	27	23	15	15
Tight roof (concrete)	2	2	2	2	3	3	2	2	3	3	2	2
Semi-tight roof (floating covers)	5	5	4	4	33	35	43	43	33	35	43	43
Natural crust	65	63	54	50	0	0	0	0	0	0	0	0
Tent, roof	28	30	40	44	37	39	40	40	37	39	40	40
Filling of storage from the bottom	95	95	95	95	90	95	95	95	75	75	75	75

Table 8.14. Detailed information on solid manure (FYM) management in animal shelters and manure storages in Finland in 2020 and as a prediction for 2025, 2030 and 2040. Unit: percentage of total solid manure per animal species.

Abatement measures	Solid manure (FYM)																							
	Cattle				Pigs				Poultry				Sheep + Goat				Horses				Fur animals			
	2020	2025	2030	2040	2020	2025	2030	2040	2020	2025	2030	2040	2020	2025	2030	2040	2020	2025	2030	2040	2020	2025	2030	2040
<u>Animal shelter</u> <u>(% of manure)</u>																								
Improved cleaning of surfaces	14	14	14	14	12	12	12	12	5	5	5	5	1	1	1	1	15	15	15	15	0	0	0	0
Flushing	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0	0	0	0
Increased manure removal frequency	4	4	4	4	2	2	2	2	95	95	95	95	0	0	0	0	-	-	-	-	0	0	0	0
Rapid urine separation	2	2	2	2	5	5	5	5	0	0	0	0	0	0	0	0	-	-	-	-	0	0	0	0
Biological or chemical air scrubbers	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	-	-	-	-	0	0	0	0
Cooling of manure channels	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Drying of manure on manure belt	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-leaking drinking system	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Manure storage</u> <u>(% of manure)</u>																								
Solid manure covering	55	60	65	65	55	60	65	65	59	60	65	65	60	60	65	65	60	60	65	65	0	0	0	0
Filling of storage from the bottom	35	35	35	35	50	50	50	50	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

Table 8.15. Detailed information on separately collected urine and dung management in animal shelters and manure storages in Finland in 2020 and as a prediction for 2025, 2030 and 2040. Unit: percentage of total urine and dung per animal species.

Abatement measures	Urine								Dung							
	Cattle				Pigs				Cattle				Pigs			
	2020	2025	2030	2040	2020	2025	2030	2040	2020	2025	2030	2040	2020	2025	2030	2040
<u>Animal shelter (% of manure)</u>																
Improved cleaning of surfaces	14	14	14	14	12	12	12	12	14	14	14	14	12	12	12	12
Flushing	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Increased manure removal frequency	-	-	-	-	-	-	-	-	4	4	4	4	2	2	2	2
Rapid urine separation	2	2	2	2	2	2	2	2	-	-	-	-	-	-	-	-
Biological or chemical air scrubbers	0	0	0	0	1	1	1	1	0	0	0	0	1	1	1	1
Cooling of manure channels	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Drying of manure on manure belt	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-leaking drinking system	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Manure storage (% of manure)</u>																
Tight roof (concrete)	8	8	8	8	8	8	8	8	-	-	-	-	-	-	-	-
Natural crust	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Tent, roof, floating cover	72	72	72	72	72	72	72	72	-	-	-	-	-	-	-	-
Solid manure covering	-	-	-	-	-	-	-	-	55	60	65	65	55	60	65	65
Filling of storage from the bottom	-	-	-	-	-	-	-	-	35	35	35	35	50	50	50	50

Table 8.16. Detailed information on slurry application in Finland in 2020 and as a prediction for 2025, 2030 and 2040. Unit: percentage of total slurry per animal species.

Abatement measures	Slurry											
	Cattle				Pigs				Poultry (no slurry systems in 2020 or later)			
	2020	2025	2030	2040	2020	2025	2030	2040	2020	2025	2030	2040
<u>Type of surface for application</u>												
Arable land	31	31	31	31	44	44	44	44	44	44	44	44
Plant covered land	44	44	44	44	31	31	31	31	31	31	31	31
Stubble	25	25	25	25	25	25	25	25	25	25	25	25
<u>Application on arable land</u>												
Injection	65	70	70	70	65	70	70	70	62	70	70	70
Band spreading	25	30	30	30	25	30	30	30	24	30	30	30
Broadcast spreading	10	0	0	0	10	0	0	0	14	0	0	0
Of manure spread on soil surface:												
No incorporation	0	0	0	0	0	0	0	0	0	0	0	0
Incorp. with ploughing < 4 h	10	15	20	20	15	15	20	20	15	15	20	20
Incorp. with ploughing < 12 h	14	14	20	20	11	14	20	20	11	14	20	20
Incorp. with ploughing 12-24 h	22	11	0	0	20	11	0	0	20	11	0	0
Incorp. with harrowing < 4 h	16	25	30	30	24	25	30	30	24	25	30	30
Incorp. with harrowing < 12 h	16	24	30	30	13	24	30	30	13	24	30	30
Incorp. with harrowing 12-24 h	22	11	0	0	17	11	0	0	17	11	0	0
<u>Application on plant covered land</u>												
Injection	65	70	70	70	65	70	70	70	62	70	70	70
Band spreading	25	30	30	30	25	30	30	30	24	30	30	30
Broadcast spreading	10	0	0	0	10	0	0	0	14	0	0	0
<u>Application on stubble</u>												
Injection	65	70	70	70	65	70	70	70	62	70	70	70
Band spreading	25	30	30	30	25	30	30	30	24	30	30	30
Broadcast spreading	10	0	0	0	10	0	0	0	14	0	0	0
Of manure spread on soil surface:												
No incorporation	0	0	0	0	0	0	0	0	0	0	0	0
Incorp. with ploughing < 4 h	10	15	20	20	15	15	20	20	15	15	20	20
Incorp. with ploughing < 12 h	14	14	20	20	11	14	20	20	11	14	20	20
Incorp. with ploughing 12-24 h	22	11	0	0	20	11	0	0	20	11	0	0
Incorp. with harrowing < 4 h	16	25	30	30	24	25	30	30	24	25	30	30
Incorp. with harrowing < 12 h	16	24	30	30	13	24	30	30	13	24	30	30
Incorp. with harrowing 12-24 h	22	11	0	0	17	11	0	0	17	11	0	0

Table 8.17. Detailed information on deep litter application in Finland in 2020 and as a prediction for 2025, 2030 and 2040. Unit: percentage of total deep litter per animal species.

Abatement measures	Deep litter														
	Cattle			Pigs			Poultry			Sheep + Goat			Horses		
	2020	2025	2030, 2040	2020	2025	2030, 2040	2020	2025	2030, 2040	2020	2025	2030, 2040	2020	2025	2030, 2040
<u>Type of surface for application</u>															
Arable land	68	68	68	88	88	88	88	88	88	68	68	68	76	76	76
Plant covered land	32	32	32	12	12	12	12	12	12	32	32	32	24	24	24
Stubble	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<u>Application on arable land</u>															
No incorporation	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Incorp. with ploughing < 4 h	15	15	20	14	15	20	14	15	20	15	15	20	14	15	20
Incorp. with ploughing < 12 h	16	25	20	13	25	20	13	25	20	16	25	20	17	25	20
Incorp. with ploughing 12-24 h	28	15	0	30	15	0	30	15	0	28	15	0	27	15	0
Incorp. with harrowing < 4 h	15	20	30	14	20	30	14	20	30	15	20	30	11	20	30
Incorp. with harrowing < 12 h	3	12	30	14	12	30	14	12	30	3	12	30	14	12	30
Incorp. with harrowing 12-24 h	23	10	0	15	10	0	15	10	0	23	10	0	17	10	0
<u>Application on plant covered land</u>															
Broadcast spreading	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100

Table 8.19. Detailed information on separately collected urine and dung application in Finland in 2020 and as a prediction for 2025, 2030 and 2040. Unit: percentage of total solid manure per animal species.

Abatement measures	Urine						Dung					
	Cattle			Pigs			Cattle			Pigs		
	2020	2025	2030, 2040	2020	2025	2030, 2040	2020	2025	2030, 2040	2020	2025	2030, 2040
<u>Type of surface for application</u>												
Arable land	17	17	17	23	23	23	68	68	68	88	88	88
Plant covered land	63	63	63	26	26	26	32	32	32	12	12	12
Stubble	20	20	20	51	51	51	-	-	-	-	-	-
<u>Application on arable land</u>												
Injection	62	70	70	62	70	70	-	-	-	-	-	-
Band spreading	24	30	30	24	30	30	-	-	-	-	-	-
Broadcast spreading	14	0	0	14	0	0	-	-	-	-	-	-
<u>Of manure spread on soil surface:</u>												
No incorporation	0	0	0	0	0	0	0	0	0	0	0	0
Incorp. with ploughing < 4 h	10	15	20	15	15	20	15	15	20	14	15	20
Incorp. with ploughing < 12 h	14	14	20	11	14	20	16	25	20	13	25	20
Incorp. with ploughing 12-24 h	22	11	0	20	11	0	28	15	0	30	15	0
Incorp. with harrowing < 4 h	16	25	30	24	25	30	15	20	30	14	20	30
Incorp. with harrowing < 12 h	16	24	30	13	24	30	3	15	30	14	15	30
Incorp. with harrowing 12-24 h	22	11	0	17	11	0	23	10	0	15	10	0
<u>Application on plant covered land</u>												
Injection	62	70	70	62	70	70	-	-	-	-	-	-
Band spreading	24	30	30	24	30	30	-	-	-	-	-	-
Broadcast spreading	14	0	0	14	0	0	100	100	100	100	100	100
<u>Application on stubble</u>												
Injection	62	70	70	62	70	70	-	-	-	-	-	-
Band spreading	24	30	30	24	30	30	-	-	-	-	-	-
Broadcast spreading	14	0	0	14	0	0	-	-	-	-	-	-
<u>Of manure spread on soil surface:</u>												
No incorporation	0	0	0	0	0	0	-	-	-	-	-	-
Incorp. with ploughing < 4 hrs	14	15	20	12	15	20	-	-	-	-	-	-
Incorp. with ploughing < 12 h	15	14	20	21	14	20	-	-	-	-	-	-
Incorp. with ploughing 12-24 h	27	11	0	22	11	0	-	-	-	-	-	-
Incorp. with harrowing < 4 h	15	25	30	15	25	30	-	-	-	-	-	-
Incorp. with harrowing < 12 h	13	24	30	14	24	30	-	-	-	-	-	-
Incorp. with harrowing 12-24 h	16	11	0	16	11	0	-	-	-	-	-	-

8.2.6. Emission reductions based on existing measures and measures that have been adopted in the legislation

This chapter presents the historic development of emissions from 2005 as well as the projections of the WM scenario up to 2030. Projections are calculated using the models described earlier in this document. The activity pathways are based on the latest Energy and Climate Strategy (prepared in 2016) which is why the estimates for 2020 are not fully in line with reported emissions of 2019. A new Strategy, complemented with a new WM scenario should be ready by the end of 2021.

Sulphur emissions

Sulphur dioxide emissions originate mainly from energy production and industrial processes. Emissions from industry decreased already between 2005-2010 in line with the limits presented in the LCPD (2001/80/EY), although the uses of both coal and peat in 2010 were higher than in 2005. Between 2010 and 2020 emissions from energy production are projected to decrease sharply due to decreasing combustion of coal, peat and HFO and the limitations in the IED. From 2020 to 2030 combustion of coal will further decrease.

Emissions from industrial processes follow the projected increase of production volumes, while a slight decrease is projected to the emission factors for metal industry and refineries due to technical improvements of processes. (Figure 8.5)




Figure 8.5. Development of SO_x emissions by sectors according to the baseline

Nitrogen oxides

The main sources for NO_x are road transport, off-road machinery and energy production. Emissions from the transport sector are projected to decrease due to EU legislation although transport volumes increase. The main contributor to decreases will be the implementation of EURO6 standards from 2015 onwards,

NO_x emissions from energy production decreased only slightly between 2005-2010 when the uses of peat, coal and biomass were restored to the normal level from their exceptional levels in 2005 when the lock-out in forest industries and the extraordinary good water situation in production of hydroelectric power decreased the demand of fuels.

The IED restricts emissions from the use of coal and biomass. The use of coal and peat also decrease notably towards 2020-2030, although biomass use is expected to increase.

Emissions from industrial processes depend on the development of production volumes and in small scale combustion on the amount of wood combusted. Impacts to emission levels from small technical improvements in both the process industry and small-scale combustion are included in the projections. (Figure 8.6)




Figure 8.6. Development of NO_x emissions by sectors according to the baseline

Particles

Important particulate matter sources are residential wood combustion, traffic, industry and peat production. In the national Energy and Climate strategy it is assumed that the combustion volume will increase slightly from 2015 to 2030, but particulate emissions will decrease due to the renewal of the combustion equipment stock.

In transport, exhaust gas emissions decrease due to the increasing number of EURO6 standard vehicles. Although direct particulate emissions in exhaust gases almost cease by 2030, traffic dust will still remain an issue. PM emissions from traffic are a significant contributor to health impacts because the emissions occur at the height of inhalation and concentrate in high density population areas.

Emissions from peat production, i.e. operations related to extraction of peat, vary annually due to peat production volumes which depend on weather (for instance between 2005–2012 from 2.7 to 5.5 kt. In the scenarios these emissions are projected to follow the projected use of peat each target year. During the last two years, peat production volumes in Finland have decreased significantly. This development is expected to continue and will probably be reflected in the renewal of the Energy and Climate Strategy.

The increasingly stringent emission regulation in combustion plants decrease emissions only slightly, since biomass consumption is expected to increase significantly

For industrial processes, no changes have been made in the projected emission factors over the years and the emissions follow development of production volumes. (Figure 8.7)




Figure 8.7. Development of small particle and black carbon emissions by sectors according to the baseline

Ammonia

The main ammonia source is agriculture where manure management drives the emissions. Small emissions are generated in transport, waste handling and industrial processes. The emission ceiling of 31 kt under the NECD and the Gothenburg Protocol is based in calculations in the RAINS model (Regional Air Pollution INformation and Simulation). In the revision of the NECD the target was to limit emissions to the level of 2010. For Finland this means a reduction of 20% in ammonia emissions from 2005, while the optimization in the GAINS would have been 15% for 2030.

In the base line approach, reductions in agricultural ammonia emissions follow the decrease in animal numbers, impacts from liquid manure systems to become more common in line with the growth of the unit size, as well as the implementation of new regulations for storage and spreading of sludge according to the updated nitrates directive (Government Decree VNa 1250/2014). On the other hand, increased production volumes raise the level of nitrogen excretion, which partly cancel the reduction by the decrease in animal numbers.

Although ammonia emissions from transport already have decreased due to improvements in technology and will further decrease, the emissions in the model are estimated at the level in 2012.

Emissions from energy production were not included in the inventory the time the FRES model was updated. These emissions will be included in the model when the inventory results are finalized. (Figure 8.8)




Figure 8.8. Development of ammonia emissions by sectors according to the baseline

NMVOG

NMVOG emissions have been decreased between 2005 - 2010 and further thereafter. The most important source is transport where emission reductions are expected due to EURO5/6 standard vehicles. Half of exhaust gas emissions originate in gasoline vehicles and half from fuel refining, storage and distribution.

LIPASTO and FRES models only covers NMVOG emission from transport and small combustion. In the WM scenario up to 2030, projections for emissions from industry and product use are based on national emission inventory values in 2016¹².

These emissions have decreased since the beginning of the 2000s' due to implementation of VOC Directives (1999/13/EC and 2004/42/EC), In Finland also the levels of activities in these sectors have decreased. For oil refineries the emission factor is estimated to decrease by 2030, however, the expected growth of the activity volume keeps the projected emission levels constant. (Figure 8.9)




Figure 8.9. Development of NMVOG emissions by sectors according to the baseline

¹² In the 2022 submission of projections, IPPU projections are updated based on estimations that are in detail explained above in Chapter Projections for IPPU and Waste (p. 31)

8.2.7 NECD 2021 review of projections

In Table 8.20 below recommendations presented in the Review Report of the 2021 NECD Review of Projections are presented with actions made on the right hand side column.

Table 8.20 - Review Report Table 4-2 Recommendations from the 2021 projections review ¹³

Observation	Year	Scenario	KC	Recommendation	Response
FI-0A-2021-0001	2021	WM	n.a	In its response to a question during the review Finland confirmed that it had accidentally copied the formula of 2020 PM2.5 into the years 2025 and 2030 in its reporting file. Finland indicated that this formula error would be corrected for its next submission. The TRT recommends that Finland include additional checks to ensure similar errors are avoided for future submissions.	The formula has been corrected.
FI-0A-2021-0003	2021	WM, WAM	n.a	For NH3 in 2020-2029, the TRT notes that under the WM scenario, Finland is predicting to miss the emissions reduction commitment. The TRT further notes that Finland did not include a WAM scenario as part of their projections submission. The TRT recommends that Finland provides for the necessary additional measures and includes them in its future projections submissions to demonstrate how all emission reduction commitments will be met.	At the moment, Finland is meeting the reduction commitment. The national air pollution control programme update in 2023 will cover updating the related projections.
FI-1A1-2019-0001	2019	WM	No	For 1A1 and 1A2 PM2.5 emissions, the TRT noted that whilst the FRES model splits 1A1 and 1A2 emissions according to the historical emissions inventory, the differences between the historical emissions inventory latest year (2018) and the 2020 projections for PM2.5 for 1A1 and 1A2 is greater than the sum of the emissions. In response to a question raised during the review, Finland explained that there was an error in the projections submissions and provided the corrected data. The TRT notes that this issue relates to an over-estimate and recommends that Finland correct the data in the next submission and ensure that appropriate quality control procedures are in place to avoid these errors.	The error has been corrected.

¹³ Where multiple pollutants are included, the issue is flagged as referring to a key category if relevant for one or more of the pollutants.

Observation	Year	Scenario	KC	Recommendation	Response
FI-1A3bvii-2019-0001	2020, 2030 2025,	With Measures (WM)	No	For 1A3bvii Automobile road abrasion, PM _{2.5} for 2020, 2025, 2030, the TRT noted that the emissions of PM _{2.5} decrease by around 15 % from 2017 to 2020. PM _{2.5} emissions are kept constant in the projection years. In response to a question raised during the review, Finland explained that no projections have earlier been made for this category (automobile road abrasion). The emissions will be revised as part of the upgraded LIPASTO system and will be included in the next reporting round. The TRT notes that this issue relates to an underestimate and recommends that revised emissions from the upgraded LIPASTO system will be included in the next emission reporting.	It was not possible to include the emissions to the 2020 submission due to need to improve the inventory methodology and changes in the organization of the inventory (all transport sector calculations were moved to VTT/Tremo). To the 2021 submission these have been included.
FI-3B-2021-0001	2021	With Measures (WM)	No	The TRT noted with reference to the whole of the agriculture sector (NFR 3) a lack of transparency regarding documentation of projected activity data and emission factors in the IIR. Also, although the external source of projected parameters was quoted in the IIR, this was not accessible to the general public. In response to a question raised during the review, Finland provided additional information on the evolution of key parameters, as well as additional explanation of the rationale for the projected parameters and references with URLs to source material. The TRT notes that this issue does not relate to an over or underestimate, and recommends that, to increase transparency, Finland include the main elements of this additional information in the projections chapter of their next IIR submission. This should include links to the publicly accessible external documentation, as well as summary-level tables or text describing the evolution of important activity data parameters (animal numbers, synthetic fertiliser application, nitrogen excretion, manure management system share, manure spreading techniques) for key categories, to make the reasons for projected trends understandable.	The IIR includes the transparent information.
FI-NATIONAL TOTAL-2019-0001	2019		n.a.	The TRT notes that Finland does not specify the historical year used as the reference year for its projections. During the review, Finland confirmed that it used 2019 from the latest (2021) inventory submission as the reference year for its projections. The TRT commends Finland for providing projections that are fully consistent with the latest historical inventory (2021 submission) despite the reporting requirement allowing projections to be consistent with the previous (2020) inventory submission. The TRT recommends that Finland clarify the year used and the inventory submission used for its reference year in future projections submissions.	The year has been inserted in the respective template cell. Finland reports here the last submission values.

Observation	Year	Scenario	KC	Recommendation	Response
FI-NATIONAL TOTAL-2019-0004	2019	With Measures (WM)	n.a.	The TRT noted that very limited information is provided in Finland's IIR on projections. The projection section provides 10 pages of summary information on trends and some limited information on methods data sources and assumptions. Although some information on its projections submitted in 2019 is available in Finland's National Air Pollution Control Programme 20305, Finland has not provided additional information or updates on methods, data sources and assumptions for projections for the 2021 submission. During the review, Finland provided some updates on methods and review of methods in response to a question from the TRT. Finland also indicated that, 'as stated in FI IIR Part 1B Table 4-1 on p. 55, the documentation will be completed to the 2022 IIR after the updating of the scenarios has been completed by the end of 2021.' The TRT recommends that Finland provides transparent information on its calculation methods, data used with references to data sources, assumptions, assessment of completeness of projections, applied PAMs/additional PAMs and sensitivity analysis (including info on parameters, variation) in its future submissions.	The previous documentation has been expanded to the 2022 submission and will be further improved when the air pollution scenarios update has been completed. The update has been postponed to 2023.

Table 8.19 - Review Report Table 4-3 Encouragements from the 2021 projections review

Scenario	Year	Scenario	KC	Encouragement	Response
FI-0A-2021-0003	2021	With Measures (WM)	No	The TRT notes that Finland expects to exceed its 2020 NH3 emission reduction commitment in its WM scenario by a small margin (2.6%) and has not presented a WAM scenario. During the review Finland provided some detailed explanation of policies and measures in place. The TRT also notes that Finland's 2030 emission reduction commitment will be achieved for NH3. The TRT encourages Finland to review its 2020 estimates and its compliance for NH3 in future submissions.	In the 2022 submission Finland complies with the ammonia reduction target.
FI-1A3biv-2021-0001	2021	With Measures (WM)	No	For 1A3biv (road transport - motorcycles) for SO2, the TRT noted that the trend for the period 2005-2020 is more than three standard deviations away from the average of the trends of all Member States and that emission estimates increased from 0.0006 kt in 2018 to 0.001 in 2020. In response to a question raised during the review, Finland explained that the projected emission estimates were reported to three decimal places and that the discrepancy was due to rounding. The TRT notes that this issue does not relate to an over or under-estimate and	The projections are reported to three decimal places in the 2022 submission. NOTE – reporting to 2 decimal places was originated from the fact that this was the default setting of the template and we

				encourages Finland to report their projected emission estimates to more decimal places in future submissions.	thought that the default should not be touched:="
FI-2A,B,C,H,I,J,K,L-2021-0001	2021	With Measures (WM)	No	For NFR 2, NMVOC, all years, the TRT noted that there was a lack of transparency regarding the allocation of NFR sectors in the projections emissions reported in Annex IV. In response to a question raised during the review, Finland explained that specific industrial processes subcategories within 2G were included in 2A,B,C,H,I,J,K,L (2D3c, 2D3d). The TRT notes that this issue does not relate to an over or under-estimate and encourages Finland to report IPPU emissions projections according to the Annex IV template.	The allocation has not yet been changed but we consider changes to the next submissions.

8.3. GRIDDED EMISSIONS AND LPS

8.3.1 Gridded data

Changes in chapter	
March 2022	TF, JM, KS
Change in methodology	New grid 2015

Background

The new EMEP grid of 0.1 degrees introduced in the 2014 Reporting Guidelines was implemented in the inventory system in 2015. Finland lies between the northern latitudes of 60° and 70°, where one degree corresponds to approximately to a 7 km *7 km area (Figure 8.10)

The presentation of gridded data in the 1° * 1° format has at the moment been implemented for the land cover of activities only in 2005. It is planned to prepare datasets also for the earlier years as well as for future years when resources are available for this kind of work.

Gridded data in the resolution of 50 km * 50 km according to the earlier versions of the Reporting Guidelines is available also for the earlier reporting years.

Submissions of gridded data are presented in Table 1.07 in IIR Part 1A.



Figure 8.10. Geographical location of Finland (Maps of the World 2016)

Developments in land use

In comparison to other European countries, Finland is still a sparsely populated country with a small urban zone in the Southern part of the country. Only the capital region is a highly urbanized area according to the classifications of EuroStat and OECD.

In addition to the low population density, a specific feature of Finland is the share of rural areas and long distances between inhabitant centres. An exceptional feature compared to other low density countries is that almost all of Finland is populated and the most distant rural areas are rather vital. In an European comparison Finland was one of the top 5 countries in the share of rural areas of total area.

During the last decades more people have moved to the population centres, rural centres of their vicinity and especially in the Southern part of Finland. Inside municipalities, population is more and more moving from sparsely populated areas to villages. Largest growth can be seen in population centres exceeding 100 000 inhabitants and secondly in 1000 - 100 000 population centres. Growth rate has been high also in centres less than 1000 inhabitants, while the sparsely populated areas continue to lose their inhabitants.

Sources included

The reporting of gridded data includes the following pollutants: NO_x (as NO₂), NMVOC, SO_x (as SO₂), NH₃, PM_{2.5}, PM₁₀, BC, CO, Pb, Cd, Hg, PCDD/F, PAH-4, HCB and PCBs. Emission data is collected from the Finnish Air Emission Information System (IPTJ). Emission and location data of installations subject to environmental permit reporting are obtained from YLVA database. Data from regional emission sources (eg. transport and agriculture) are based on calculation and have been geographically distributed on the basis of more detailed national emission data.

The emission source classifications are based on the UN classification of climate and long-range transport agreements and GNFR sectors reported are A_PublicPower, D_Fugitive, B_Industry, C_OtherStationaryComb, I_Offroad, H_Aviation, F_RoadTransport, G_Shipping, E_Solvents, M_Other, J_Waste, K_AgriLivestock and L_AgriOther. (Tables 8.21-8.22)

Table 8.21 – Air pollutant emissions of GNFR categories for the year 2018.

Pollutant	Unit	A_PublicPower	B_Industry	C_OtherStationaryComb	D_Fugitive	E_Solvents	F_RoadTransport	G_Shipping	H_Aviation	I_Offroad	J_Waste	K_AgriLivestock	L_AgriOther
NOx	t	24357.9	29347.2	10098.6		6.1	30429.7	6472.7	1017.8	15420.5		423.5	9020.9
NMVOC	t	1603.3	15041.4	22130.0	6075.5	9860.3	5111.1	2937.2	140.2	6171.5	88.4	12978.6	3161.7
SOx	t	12457.4	16699.1	3667.4	52.8	5.0	46.8	84.4	64.1	42.2			7.9
NH3	t	3.7	659.9	1172.6	3.2	234.1	838.1	1.0		6.3	475.1	18324.0	10471.5
PM2.5	t	336.3	2770.2	9541.2	1189.5	203.3	1911.7	298.4	7.4	919.6	103.9	166.5	350.0
PM10	t	1178.1	4140.0	10574.6	1697.0	214.8	7560.0	330.4	7.4	924.8	104.4	571.4	3812.8
BC	t	18.0	75.3	2704.8	0.0	3.7	704.0	61.4	3.7	414.4	9.4		18.8
CO	t	15646.9	33040.7	167383.1		187.1	37153.7	19565.3	1204.9	74447.8			1901.5
Pb	kg	2188.9	9983.0	1573.7	4.0	1113.3	491.7	12.6	36.3	3.7	1.2		1.6
Cd	kg	134.4	458.0	244.2	0.2	19.4	2.2	1.0		8.4	0.7		14.6
Hg	kg	160.0	424.7	42.8	0.0	0.1	25.1	2.7		0.9	18.1		2.9
PCDD/F	g	3.7	4.4	1.6	2.6	0.0	1.0	0.0		0.0	1.1		0.0
PAHs	g	546245.6	366667.6	8318764.0	456176.8	11384.1	225747.3			64521.5	1000.0		10.6

HCB	kg	0.5	31.0	0.3		0.0	0.2	0.0		0.0	0.0		0.0
PCBs	g	319.7	19145.1	3760.3	3098.6		0.2	9.2		1.1	12.1		

The categories contain point sources and non-point sources as illustrated in the table below.

Table 8.21 - The relative shares of emissions from point and non-point sources per aggregate category

Pollutant	Public power and industries*		Other stationary combustion**		Traffic and agriculture***		Products and waste****	
	Point sources	Non-point sources	Point sources	Non-point sources	Point sources	Non-point sources	Point sources	Non-point sources
NOx	98.5 %	1.5 %	0.4 %	99.6 %	0.0 %	100.0 %	0.0 %	100.0 %
NMVOc	50.8 %	49.2 %	0.0 %	100.0 %	0.0 %	100.0 %	21.0 %	79.0 %
SOx	98.1 %	1.9 %	0.3 %	99.7 %	0.0 %	100.0 %	14.1 %	85.9 %
NH3	99.5 %	0.5 %	0.0 %	100.0 %	0.0 %	100.0 %	31.2 %	68.8 %
TSP	58.8 %	41.2 %	0.1 %	99.9 %	0.0 %	100.0 %	20.1 %	79.9 %
PM10	63.3 %	36.7 %	0.0 %	100.0 %	0.0 %	100.0 %	20.2 %	79.8 %
PM2.5	62.7 %	37.3 %	0.0 %	100.0 %	0.0 %	100.0 %	19.0 %	81.0 %
BC	98.6 %	1.4 %	0.0 %	100.0 %	0.0 %	100.0 %	0.0 %	100.0 %
CO	99.2 %	0.8 %	0.0 %	100.0 %	0.0 %	100.0 %	0.0 %	100.0 %
Pb	94.4 %	5.6 %	0.3 %	99.7 %	0.0 %	100.0 %	0.0 %	100.0 %
Cd	98.1 %	1.9 %	0.0 %	100.0 %	0.0 %	100.0 %	0.0 %	100.0 %
Hg	98.2 %	1.8 %	0.2 %	99.8 %	0.0 %	100.0 %	0.0 %	100.0 %
As	84.4 %	15.6 %	0.8 %	99.2 %	0.0 %	100.0 %	0.0 %	100.0 %
Cr	94.9 %	5.1 %	0.1 %	99.9 %	0.0 %	100.0 %	0.0 %	100.0 %
Cu	85.2 %	14.8 %	0.4 %	99.6 %	0.0 %	100.0 %	0.0 %	100.0 %
Ni	94.1 %	5.9 %	0.1 %	99.9 %	0.0 %	100.0 %	0.0 %	100.0 %
Zn	97.1 %	2.9 %	0.0 %	100.0 %	0.0 %	100.0 %	0.0 %	100.0 %
PCCD/F	98.0 %	2.0 %	0.1 %	99.9 %	0.0 %	100.0 %	0.0 %	100.0 %
PAH-4	99.0 %	1.0 %	0.0 %	100.0 %	0.0 %	100.0 %	13.0 %	87.0 %
HCB	99.3 %	0.7 %	0.0 %	100.0 %	0.0 %	100.0 %	0.0 %	100.0 %
PCB	85.3 %	14.7 %	0.0 %	100.0 %	0.0 %	100.0 %	0.0 %	100.0 %

The headers aggregate the GNFR categories as follows:

* Public power and industries: A_PublicPower, D_Fugitive, B_Industry

** Other stationary combustion: C_OtherStationaryComb

*** Traffic and agriculture: I_Offroad, H_Aviation, F_RoadTransport, G_Shipping, K_AgriLivestock, L_AgriOther

**** Products and waste: E_Solvents, M_Other, J_Waste

Point sources are distributed by Tier 3 methodology. IPTJ contains coordinate data as WGS84 for all known point sources. Non-point sources vary between Tier 1 and 2 based on the activity. The table below presents the primary tier and the secondary tier in brackets. The secondary tier applies to the point or non-point emissions of the category which contributes to the lesser half of total emissions (for categories with both emission sources present). (Table 8.22)

Table 8.22 – Tier categorisation of spatial disaggregation of GNFR-categories

GNFR19	A_PublicPower	B_Industry	C_OtherStationaryComb	D_Fugitive	E_Solvents	F_RoadTransp	G_Shipping	H_Aviation	I_Offroad	J_Waste	K_AgriLivestoc	L_AgriOther
Tier	3 (1)	3 (1)	1 (3)	1	1 (3)	2	2	2	2	1 (3)	1	1

Data sources for disaggregation of non-point emission sources

Changes in chapter	
March 2022	JM

The geographical distribution of non-point emission sources, such as transport, households, agriculture and small-scale wood burning utilizes Finnish-wide proxies, which aim to represent each emission source with highest applicable level of accuracy. The spatial data set of the national road and street information system (Digiroad) is used as a medium for traffic emissions. Numerous different data sources have been utilized in allocating the calculated regional emissions to the map. The most commonly used source material is Corine Land Cover (CLC2006), which describes the Finnish land use and land cover in 2006. The material covers built land, agricultural areas, forests, open canals and rocky lands, wetlands and bogs and water areas. The data extracted from CLC2006 is supplemented with SLICES (Separated Land Use & Cover information System) data. Built environment is modelled with the Building and Housing Register (RHR) data. Activities without suitable proxies are distributed according to population density.

Point source sources, such as power plants and industrial operational plants, are shown according to their coordinates. Non-point emission sources such as emissions from transport, consumption and production, agriculture, and small-scale wood burning, cannot be allocated to a single point, but will use the indirect data that best represent each emission source. An activity for which no suitable medium exists is distributed according to population density. The proxies are currently based on land use of 2010 +/- 5 years depending on the availability of the data.

The proxies for non-point emission-sources are linked to the emissions by SNAP categorization. The table below presents the proxies and their corresponding SNAP, NFR and GNFR categorizations as are used in the GRID inventory submission of 2020 (Table 8.23).

Table 8.23 – List of used proxies per SNAP category in the inventory submission of 2020 of gridded emissions

GNFR19	NFR	NFR Description	SNAP	SNAP Description	Proxy	Source
A_Public Power	1A1a	Public electricity and heat production	010205	District heating - Stationary engines	Industrial areas and services	Corine land cover 2006
B_Industry	2C1	Iron and steel production	040208	Rolling mills	Industrial areas and services	Corine land cover 2006
	2H2	Food and beverages industry	040627	Meat, fish etc. frying / curing	Industrial areas and services	Corine land cover 2006
	2B1	Ammonia production	040403	Ammonia	Industrial areas and services	Corine land cover 2006
	2D3c	Asphalt roofing	040610	Asphalt roofing materials	All buildings - floor area	National building and dwelling register
	2I	Wood processing	040620	Wood processing	Industrial areas and services	Corine land cover 2006
	2C7c	Other metal production	040309z	Other	Industrial areas and services	Corine land cover 2006
	2H2	Food and beverages industry	040606	Wine	Wineries	Public listing
	2A5b	Construction and demolition	040624	Public works and building sites	All buildings - floor area	National building and dwelling register
	2H2	Food and beverages industry	040625	Sugar production	Industrial areas and services	Corine land cover 2006
	2A5c	Storage, handling and transport of mineral products	040900	Storage, handling and transport of mineral products	Industrial areas and services	Corine land cover 2006

2B10a	Chemical industry: Other (Please specify in the IIR)	040407	NPK fertilisers	Industrial areas and services	Corine land cover 2006
2B10b	Storage, handling and transport of chemical products (Please specify in the IIR)	040415	Storage and handling of inorganic chemical prod. (o)	Mines and industrial areas	Separated Land Use/Land Cover Information System (SLICES)
2H2	Food and beverages industry	040605	Bread	Industrial areas and services	Corine land cover 2006
2H2	Food and beverages industry	040607	Beer	Breweries	Public listing
2A5a	Quarrying and mining of minerals other than coal	040616	Extraction of mineral ores	Mines	Separated Land Use/Land Cover Information System (SLICES)
1A2gviii	Stationary combustion in manufacturing industries and construction: Other (Please specify in the IIR)	030326	Other	Facilities for energy production	National building and dwelling register
2A3	Glass production	040613	Glass (decarbonizing)	Industrial areas and services	Corine land cover 2006
2C7c	Other metal production (Please specify the sources included/excluded in the notes column to the right)	040210	Other	Industrial areas and services	Corine land cover 2006
2C3	Aluminium production	040301	Aluminium production (electrolysis)	Industrial areas and services	Corine land cover 2006
2H1	Pulp and paper industry	040602	Paper pulp (kraft process)	Industrial areas and services	Corine land cover 2006
2L	Other production, consumption, storage, transportation or handling of bulk products	040617	Other (including asbestos products manufacturing)	Industrial areas and services	Corine land cover 2006
2D3b	Road paving with asphalt	040611	Road paving with Asphalt	Streets under construction	Digiroad
2C7d	Storage, handling and transport of metal products	040211	Storage, handling and transport of ferrous metal products	Population density	National building and dwelling register
2A2	Lime production	040614	Lime (decarbonizing)	Industrial areas and services	Corine land cover 2006
2C1	Iron and steel production	040209	Sinter and pelletizing plant (except comb. 030301)	Industrial areas and services	Corine land cover 2006
2A1	Cement production	040612	Cement (decarbonizing)	Industrial areas and services	Corine land cover 2006
C_Other Stationary Comb	1A5a Other stationary (including military)	020106	Commercial and institutional - Other stationary equipment (n)	Industrial areas and services	Corine land cover 2006
	1A4ci Agriculture/Forestry/Fis hing: Stationary	020305	Agri./forest/aqua. - Other stationary equipment (n)	Built agricultural land	Separated Land Use/Land Cover

						Information System (SLICES)
	1A4bi	Residential: Stationary	020205	Residential - Other equipment (stoves, fireplaces, cooking)	Buildings with wood as the primary heat source	National building and dwelling register
	1A4ai	Commercial / institutional: Stationary	020103b	Commercial and institutional - Combustion plants < 20 MW (boilers)	Population density	National building and dwelling register
	1A5a	Other stationary (including military)	020103b	Commercial and institutional - Combustion plants < 20 MW (boilers)	Population density	National building and dwelling register
	1A4bi	Residential: Stationary	020202b	Residential - Combustion plants < 20 MW (boilers)	Population density	National building and dwelling register
	1A4ai	Commercial / institutional: Stationary	020106	Commercial and institutional - Other stationary equipment (n)	Industrial areas and services	Corine land cover 2006
	1A4ci	Agriculture/Forestry/Fishing: Stationary	020302b	Agri./forest/aqua. - Combustion plants < 20 MW (boilers)	Population density	National building and dwelling register
D_Fugitive	1B2b	Fugitive emissions from natural gas (exploration, production, processing, transmission, storage, distribution and other)	050601	Pipelines (q)	Industrial areas and services	Corine land cover 2006
	1B1c	Other fugitive emissions from solid fuels	050121	Peat production	Population density	National building and dwelling register
	1B2av	Distribution of oil products	050502	Transport and depots (except 050503)	Industrial areas and services	Corine land cover 2006
	1B1b	Fugitive emission from solid fuels: Solid fuel transformation	040201	Coke oven (door leakage and extinction)	Industrial areas and services	Corine land cover 2006
	1B2av	Distribution of oil products	050503	Service stations (including refuelling of cars)	Service stations	National building and dwelling register
E_Solvents	2G	Other product use	060601	Use of fireworks	Population density	National building and dwelling register
	2D3e	Degreasing	060204	Other industrial cleaning	All buildings - volume	National building and dwelling register
	2D3g	Chemical products	060314	Other	Population density	National building and dwelling register
	2D3i	Other solvent use	060412	Other (preservation of seeds,...)	Agricultural land in use (overall)	Separated Land Use/Land Cover Information System (SLICES)
	2D3i	Other solvent use	060406	Preservation of wood	All buildings - floor area	National building and dwelling register
	2D3a	Domestic solvent use including fungicides	060408	Domestic solvent use (other than paint application)	Buildings used for permanent residence	National building and dwelling register

	2D3d	Coating applications	060103	Paint application : construction and buildings (except item 060107)	All buildings - floor area	National building and dwelling register
	2D3d	Coating applications	060108	Other industrial paint application	Industrial areas and services	Corine land cover 2006
	2D3g	Chemical products	060310	Asphalt blowing	Population density	National building and dwelling register
	2D3d	Coating applications	060109	Other non-industrial paint application	All buildings - floor area	National building and dwelling register
	2D3h	Printing	060403	Printing industry	Population density	National building and dwelling register
	2D3i	Other solvent use	060404	Fat, edible and non-edible oil extraction	Population density	National building and dwelling register
	2D3g	Chemical products	060313	Leather tanning	Population density	National building and dwelling register
	2D3g	Chemical products	060307	Paints manufacturing	Population density	National building and dwelling register
	2G	Other product use	060602	Use of tobacco	Population density	National building and dwelling register
F_Road Transport	1A3bv	Road transport: Gasoline evaporation	070600	Gasoline evaporation from vehicles	Service stations	National building and dwelling register
	1A3biii	Road transport: Heavy duty vehicles and buses	070300	Heavy duty vehicles > 3.5 t and buses	Streets and roads (weighed with no. vehicles)	Digiroad
	1A3biv	Road transport: Mopeds & motorcycles	070500	Motorcycles > 50 cm3	Streets and roads (weighed with no. vehicles)	Digiroad
	1A3bi	Road transport: Passenger cars	070100	Passenger cars	Streets and roads (weighed with no. vehicles)	Digiroad
	1A3bvii	Road transport: Automobile road abrasion	070800	Road abrasion	Streets and roads (weighed with no. vehicles)	Digiroad
	1A3bii	Road transport: Light duty vehicles	070200	Light duty vehicles < 3.5 t	Streets and roads (weighed with no. vehicles)	Digiroad
	1A3bvi	Road transport: Automobile tyre and brake wear	070700	Automobile tyre and brake wear	Streets and roads (weighed with no. vehicles)	Digiroad
G_Shipping	1A3dii	National navigation (Shipping)	080303	Personal watercraft	Water bodies over 200 hectares and built water environments	Other
	1A3dii	National navigation (Shipping)	080304	Inland goods carrying vessels	Water bodies over 200 hectares and built water environments	Other
	1A3dii	National navigation (Shipping)	080302	Motorboats / workboats	Water bodies over 200 hectares and built water environments	Other
	1A3dii	National navigation (Shipping)	080402	National sea traffic within EMEP area	National ports (incl. onland area occupied)	Corine land cover 2006
H_Aviation	1A3ai(i)	International aviation LTO (Civil)	080502	International airport traffic (LTO cycles - <1000 m)	Airports (weighed with passenger numbers)	Separated Land Use/Land Cover Information System (SLICES)

	1A3aii(i)	Domestic aviation LTO (Civil)	080501	Domestic airport traffic (LTO cycles - <1000 m)	Airports (weighed with passenger numbers)	Separated Land Use/Land Cover Information System (SLICES)
I_Offroad	1A3c	Railways	080200	Railways	Railroads	Railroads
	1A4ciii	Agriculture/Forestry/Fishing: National fishing	080403	National fishing	Water bodies over 200 hectares and built water environments	Other
	1A4aii	Commercial / institutional: Mobile	081000	Other off-road	All buildings - count	National building and dwelling register
	1A5b	Other, Mobile (including military, land based and recreational boats)	080100	Military	Population density	National building and dwelling register
	1A4cii	Agriculture/Forestry/Fishing: Off-road vehicles and other machinery	080700	Forestry	New tree stumps from 2009 to 2011	Finnish Forest Institute
	1A4cii	Agriculture/Forestry/Fishing: Off-road vehicles and other machinery	080600	Agriculture	Agricultural land in use (overall)	Separated Land Use/Land Cover Information System (SLICES)
	1A2gvii	Mobile Combustion in manufacturing industries and construction: (Please specify in the IIR)	080800	Industry – off road	Industrial areas and services	Corine land cover 2006
	1A4bii	Residential: Household and gardening (mobile)	080900	Household and gardening	Buildings used for permanent residence	National building and dwelling register
	J_Waste	5D1	Domestic wastewater handling	091002	Waste water treatment in residential and commercial sect.	Population density
5C1bv		Cremation	090901	Incineration of corpses	Crematoriums	Public listing
5A		Biological treatment of waste - Solid waste disposal on land	090401	Managed Waste Disposal on Land	Landfills	Other
5C1biii		Clinical waste incineration	090207	Incineration of hospital wastes	Population density	National building and dwelling register
5B1		Biological treatment of waste - Composting	091005	Compost production	Sparse residential areas	Corine land cover 2006
5E		Other waste (Please specify in IIR)	091101	Unintentional house fires	Population density	National building and dwelling register
5A		Biological treatment of waste - Solid waste disposal on land	090403	Other	Population density	National building and dwelling register
5E		Other waste (Please specify in IIR)	091007	Latrines	Sparse residential areas	Corine land cover 2006
5D2		Industrial wastewater handling	091001	Wastewater treatment in industry	Industrial areas and services	Corine land cover 2006
5E		Other waste (Please specify in IIR)	091102	Unintentional car fires	Population density	National building and dwelling register
K_Agri Livestock	3B4giii	Manure management - Turkeys	100509a	Turkeys	Farmhouses (Swine and bovine)	National building and dwelling register
	3B1a	Manure management - Dairy cattle	100501	Dairy cows	Farmhouses (Swine and bovine)	National building and dwelling register

	3B3	Manure management - Swine	100504	Sows	Farmhouses (Swine and bovine)	National building and dwelling register
	3B2	Manure management - Sheep	100505	Sheep	Farmhouses (Other animals)	Separated Land Use/Land Cover Information System (SLICES)
	3B4gi	Manure management - Laying hens	100507	Laying hens	Farmhouses (Swine and bovine)	National building and dwelling register
	3Da3	Urine and dung deposited by grazing animals	100517	Urine and dung deposited by grazing animals	Fields and fallows	Separated Land Use/Land Cover Information System (SLICES)
	3B4e	Manure management - Horses	100506	Horses	Stables	National building and dwelling register
	3B4gii	Manure management - Broilers	100508	Broilers	Farmhouses (Other animals)	Separated Land Use/Land Cover Information System (SLICES)
	3B4giv	Manure management - Other poultry (please specify in the IIR)	100509z	Other poultry (ducks, geese ,etc.)	Population density	National building and dwelling register
	3B4h	Manure management - Other animals (please specify in the IIR)	100510	Fur animals	Farmhouses (Other animals)	Separated Land Use/Land Cover Information System (SLICES)
	3B4h	Manure management - Other animals (please specify in the IIR)	100516	Reindeer	Population density	National building and dwelling register
	3B1b	Manure management - Non-dairy cattle	100502	Other cattle	Farmhouses (Swine and bovine)	National building and dwelling register
	3B3	Manure management - Swine	100503	Fattening pigs	Farmhouses (Swine and bovine)	National building and dwelling register
	3B4d	Manure management - Goats	100511	Goats	Farmhouses (Other animals)	Separated Land Use/Land Cover Information System (SLICES)
L_AgriOther	3Da1	Inorganic N-fertilizers (includes also urea application)	100104	Market gardening	Fields and fallows	Separated Land Use/Land Cover Information System (SLICES)
	3Df	Use of pesticides	100600	Use of pesticides and Limestone	Fields and fallows	Separated Land Use/Land Cover Information System (SLICES)
	3Da2b	Sewage sludge applied to soils	100906	Sewage sludge applied to soils	Fields and fallows	Separated Land Use/Land Cover Information System (SLICES)
	3Dc	Farm-level agricultural operations including storage, handling and transport of agricultural products	101000	Farm-level storage, handling and transport of agricultural products	Built agricultural land	Separated Land Use/Land Cover Information System (SLICES)
	3Da1	Inorganic N-fertilizers (includes also urea application)	100101	Permanent crops	Fields and fallows	Separated Land Use/Land Cover Information System (SLICES)

	3Da2a	Animal manure applied to soils	100905	Animal manure applied to soils	Fields and fallows	Separated Land Use/Land Cover Information System (SLICES)
	3F	Field burning of agricultural residues	100300	On-field burning of stubble, straw, etc.	Fields and fallows	Separated Land Use/Land Cover Information System (SLICES)
	3Da1	Inorganic N-fertilizers (includes also urea application)	100102	Arable land crops	Fields and fallows	Separated Land Use/Land Cover Information System (SLICES)
	3Db	Indirect emissions from managed soils	100208	Indirect emissions from managed soils	Fields and fallows	Separated Land Use/Land Cover Information System (SLICES)
	3Da1	Inorganic N-fertilizers (includes also urea application)	100105	Grassland	Fields and fallows	Separated Land Use/Land Cover Information System (SLICES)
O_AviCruise	1A3ai(ii)	International aviation cruise (Civil)	080504	International cruise traffic (>1000 m)	Population density	National building and dwelling register
	1A3aii(ii)	Domestic aviation cruise (Civil)	080503	National cruise traffic (>1000 m)	Population density	National building and dwelling register
P_IntShippi ng	1A3di(i)	International maritime navigation	080404	International sea traffic (international bunkers)	Population density	National building and dwelling register

Corine Land Cover 2006

CORINE Land Cover 2006 (CLC2006) dataset provides information on Finnish land cover and land use. The data is derived from the European CLC 2006 project and it includes raster data with the resolution of 25 x 25 metres. The data is produced by SYKE based on automated interpretation of satellite images and data integration. The standard CLC nomenclature contains 44 categories for land cover of which the following are selected to be used as basis of distribution:

- Class 1110 — Dense residential
- Class 1120 — Sparse residential
- Class 1210 — Industry and Services
- Class 1220 — Transportation
- Class 1230 — Harbour areas
- Class 1310 — Land extraction areas
- Class 1320 — Landfills
- Class 1330 — Construction areas
- Class 1421 — Second houses and non-permanent living

The data is extracted from original source and converted from raster data into point. These points are aggregated into EMEP grid cells, where their total count within a cell acts as the density defining factor for the given cell. The accuracy of the method is dependent on the accuracy of the source material. Using land use as the basis for basis for distribution of diffuse emissions does not take into account the rate of activity within the given area. Some accuracy is also lost during conversion. However, an example analysis made for land extraction areas reveals a correlation with land use and the Salpausselkä ridge system. which is known of possessing a high activity rate for land extraction. (Figure 8.11)




Figure 8.11 – CLC2006 marked land extraction areas in EMEP grid (background) and their correlation with the Salpausselkä ridge system (foreground)

SLICES

Separated Land Use/Land Cover Information System (SLICES) is a land use dataset from a joint operation between National Land Survey of Finland (NLS), Finnish Environment Institute (SYKE) and the Finnish Forest Research Institute (METLA, currently known as the Natural Resources Institute Finland). As a source material it is handled with the same principles as CLC data, as SLICES is a sub-constituent to CLC material. The categorization of data however differs and the following models are extracted based on SLICES data:

- Airports (by land use)
- Storage areas
- Agricultural areas with activity
- Agricultural fallows
- Built agricultural land
- Mines and other mineral extraction
- Extraction of sand, gravel and other land extraction

The emissions of GNFR H_Aviation are distributed to the land used by airports, which are weighed by the statistics of Finavia for the year 2013. Table 8.24 presents the passenger numbers for selected cities.

Table 8.24 – Passenger numbers per airport used to scale aviation emissions

ObjectID	Municipality	Passengers
32	Vantaa	15278994
190	Oulu	877080
204	Pirkkala	466671
234	Rovaniemi	427367
284	Turku	324687
296	Vaasa	319315
252	Siilinjärvi	261151
97	Kittilä	237222
49	Inari	146314
146	Liperi	131291
31	Helsinki	100000
ObjectID	Municipality	Passengers
133	Lappeenranta	98300
119	Kuusamo	74583
71	Kajaani	74558

111	Kruunupyy	68991
88	Kemi	57681
65	Jyväskylä	50570
207	Pori	26229
11	Enontekiö	20169
246	Savonlinna	12215
59	Joroinen	6759
109	Kouvola	500

For some airports no passenger data is available. For these 100 passengers is assumed. These municipalities are Alajärvi, Alavus, Asikkala, Imatra, Jomala, Jämijärvi, Jämsä, Kauhajoki, Kauhava, Keminmaa, Kitee, Kokemäki, Kontiolahti, Kotka, Kuhmo, Kemijärvi, Lieksa, Loppi, Oripää, Pudasjärvi, Raahe, Rautavaara, Salo, Sodankylä, Suomussalmi, Tampere, Vihti, Ylivieska, Eura. Hyvinkää airport is assumed to be used only for non-motorized air vehicles.

Building and Dwelling Register

Building and Dwelling Register (here referred as BDR, fin Rakennus- ja Huoneistorekisteri. RHR) is the national register for buildings and dwellings. The use of the register is limited by several laws. The data of BDR is used only as the basis of analysis. No detailed information or exact counts of buildings or housing in an area can by any means be backtracked based on the published data. Based on BDR data the following models are constructed to be used as basis for distribution:

- Population density
- All buildings – by count
- All buildings – by floor area
- All buildings – by overall volume
- Residential buildings – permanent
- Residential buildings – temporary
- Energy production facilities
- Energy production facilities with wood based fuels as primary heat source
- Commercial buildings with wood based fuels as primary heat source
- Residential buildings with wood based fuels as primary heat source
- Agriculture – Horse stables and other animal shelters
- Agriculture – Piggeries, cattle shelters and henhouses
- Agriculture – grain drying kilns and facilities
- Petrol stations and other automobile service facilities

For models based on all buildings several models were constructed based on count, floor area and the overall volume to be used for different purposes. For example for categories of product use the it can be assumed that the activity rate is more connected to the count of households rather than volumetric parameters. For emissions resulting from heating, the overall volume gives a better estimate as the amount of energy required correlates with building volume. The inventory of 2022 will substitute the previous model with a more detailed model, both of which are based on the data above.

National road and street information system (Digiroad)

Digiroad is the national road- and street information system which contains the centre line geometry data for all known streets. The activity rate is also available for most main roads as vehicles per day. The following activities are extracted for use as proxies:

- Highways
- Roads
- Streets
- Private streets
- All combined

Activity rate is introduced with the following formula:

$$E_{cell} = \frac{E_{total}}{n_{total}} * n_{cell}$$

where

E_{cell} = Emission value in a cell

E_{total} =Sum value of emissions

n_{total} =Vehicles per day (total national)

n_{cell} =Vehicles per day (cell)





Figure 8.12 illustrates the resulting map after applying the methodology.

Figure 8.12 - PM_{2.5} emissions of 2013 in the national street system

Other sources

In the Finnish emission inventory crematoria, small breweries and wineries are calculated as diffuse sources. It is however possible to collect the location data for most of these based on various public listings. The data extracted in this manner is geocoded based on publicly available addresses and emissions are assigned to the corresponding cells. The method does not take the rate of activity within the locations into account.

For agricultural emissions a pre-existing proxy containing information about field distribution is used. This proxy is used for GNFR L_AgriOther under which Finland reports the following activities: indirect emissions from managed soils, urine and dung deposited by grazing animals, inorganic N-fertilizers, on-field burning of stubble, use of pesticides, animal manure applied to soils, sewage sludge applied to soils and farm-level agricultural operations. Spatially these emissions are estimated by fields and built agricultural land. Agricultural activities concentrate on the south-west of Finland and there is a notable amount of fields in the vicinity of the largest cities. To illustrate further, the images below present an image of the distribution of emissions before and after reconfiguring the proxies for the 2020 submission as a response to observation FI-GRID-L-2020-0001. Figure 8.14 presents the registered fields for Southern Finland (as listed by the Finnish Food Authority).



Initial submission Revised estimate



Registered fields in the registry of Finnish food authority

Figure 8.13. Maps above: Distribution of emissions before and after reconfiguring the proxies for the 2020 submission as a response to observation FI-GRID-L-2020-0001. Map below: Registered fields for Southern Finland (as listed by the Finnish Food Authority) marked in pink colour.

Public viewing tool for spatial distribution of emissions

A viewing tool of the spatial distribution of emissions within the inventory can be found at: <https://www.p2.ymparisto.fi/paastotkartalla/?lang=en>. The viewing tool aggregates GNFR sectors as presented in Table 8.25.

Table 8.25 – Description of the aggregation of GNFR classes for public viewing.

GNFR	Aggregate class	Description of sources
A_PublicPower	Public power and industries	Energy production and industrial processes covers emissions from electricity production and district heating, manufacturing and handling of fuels as well as from industrial processes and industrial boilers. Emissions which can not be allocated to geographical locations are evenly distributed on the map on industrial areas.
D_Fugitive		
B_Industry		
C_OtherStationaryComb	Other stationary combustion	Other stationary combustion includes such small scale furnaces and heat sources. Most of these are heat sources for housing, but also included are sauna stoves, fireplaces and stoves using wood or biomass as the primary fuel. The category also includes emissions from known fuel consumption that can not be allocated to a known facility.
I_Offroad	Transport	Transport category includes emissions from road and rail transport, navigation, aviation, non-road and working machinery. Regarding road transport emissions from tyre, brake wear and road abrasion as well as gasoline evaporation are included. The mapping has been carried out using data on road network, starting points of navigation and aviation without route data.
H_Aviation		
F_RoadTransport		
G_Shipping		
E_Solvents	Products and waste	Product use and waste includes use of products and solvents in industry and households. Emissions from landfill, composting and wastewater treatment are included, as well as e.g. house and car fires.
M_Other		
J_Waste		
K_AgriLivestock	Agriculture	The emissions from agriculture consists of such activities as animal husbandry and manure management, fertilizing and the use of pesticides. Also field burning of agricultural residues and other field operations are listed under agricultural emissions.
L_AgriOther		

Methodological issues

Changes in chapter	
March 2021	JM

Overview of the calculation of emissions

Gridded emissions contain all of the emissions of the air emission inventory including point sources and non-point sources. The national emission values are distributed to the cells with the following generalized equation.

$$E_{cell} = \frac{E_{total}}{n_{total}} * n_{cell}$$

Where

- E_{cell} = Emission value in a cell
- E_{total} = Total emission value
- n_{total} = Rate of activity, national total
- n_{cell} = Rate of activity in a cell

The methodology is constructed by the instructions of *Spatial mapping of emissions* of EMEP/EEA Guidebook 2013. As of the latest methodology of Guidebook 2019, the following steps are included in the methodology:

- Key category analysis to identify the most important sources is used.
- Existing spatial datasets are preferred.
- GIS tools are used to improve the proxies.
- Proxy data that is judged to most closely represent the spatial emissions patterns and intensity, and which is applicable with available resources, is selected.
- Spatial datasets that are complete are preferred.
- New data is available rarely. The currently gridded data uses spatial proxies which are set to describe land use of the year 2010.
- Issues relating to non-disclosure may be encountered but have not been observed to date.
- Aggregation is done in the EMEP 0.1 x 0.1 degree longitude/latitude grid but the methodology of creating proxies is applicable to higher levels of detail when needed

Disaggregating diffuse emissions.

The methodology to create proxies follows the basic principles as presented in the Guidebook 2019 where applicable, with modifications based on source content. Point sources are gridded based on chapter 3.4.1. Area sources are gridded based on chapter 3.4.2, however in many cases the material is first converted into points of certain density within the area. This commonly takes place with rasterized data sources that do not allow spatial analysis with the available toolset. Line sources are gridded according to chapter 3.4.3. Converting spatial projections is done where needed.

Uncertainty and time series' consistency

No evaluation of uncertainty has been done for gridded data. The methodology is prone to uncertainty especially in the temporal scale, as spatial disaggregation is done only for the target year 2010.

Source-specific QA/QC and verification

Normal statistical quality checking related to assessment of magnitude and trends has been carried out. Visual inspection for all GNFR sectors.

Source-specific recalculations including changes made in response to the review process

2020

- Update of methodology and inclusion of new inspection tools.
- Correction of geographical allocation issues: FI-GRID-L-2020-0001, FI-GRID-C-2020-0001, FI-GRID-B-2020-0001
- Update of coordinates to selected power plants (in response to FI-GRID-A-2020-0001)

2021

- Rewritten documentation of gridded emissions (in response to *Notes on reporting of air pollutant emissions from Large Point sources and emissions gridded data under the NECD*)
- Improved accuracy in facility coordinates
- Facility locations unified with LPS

Source-specific planned improvements

Substituting the proxy for small scale wood combustion to the more accurate FRES model (Finnish Regional Emission Scenario). Scheduled spring 2022.

8.3.2 LPS data, sources, geographical coordinates and emissions

Changes in chapter	
March 2022	TF
Change in method	none

According to the 2014 CLRTAP Reporting Guidelines (ECE/EB.AIR/125), large point sources (LPS) are defined as facilities whose combined emissions, within the limited identifiable area of the site premises, exceed the pollutant emission thresholds identified in table 1 of the reporting guidelines. In the Finnish inventory, LPS facilities are identified as facilities that have at least one E-PRTR installation at their site premises. All emissions from all installations of these facilities, that are located at their site premises and exceed the pollutant emission thresholds identified in the reporting guidelines, are included in the LPS reporting data.

The emission data from LPS facilities are either reported by the facilities according to the environmental monitoring requirements in their environmental permits, and available from the YLVA system, or calculated at SYKE. Always when facility reported emission data are available, it is used in the reporting. All the reported LPS emission data is also included in the national inventory.

The emission data reported by Finland under the E-PRTR regulation is extracted from the YLVA system. This emission data is also included in the national inventory and LPS emission data. The differences in the emission data of individual facilities reported under the E-PRTR and LPS reporting are usually due to the inclusion of all installations from the site premises of the facilities to the LPS reporting. The E-PRTR reporting comprise only E-PRTR installations. Furthermore, in some cases, the LPS reporting data may also contain emissions calculated at SYKE in addition to the emissions reported by the facility to the YLVA system. Sometimes, erroneous emission data of E-PRTR installations is detected from the YLVA system (e.g. emissions reported in a wrong unit). When detected, these are corrected to the inventory data extracted from YLVA. The facility supervisors from the Centres for Economic Development, Transport and the Environment are informed of these data. If these data are not corrected to the E-PRTR reporting, this causes also differences between inventory/LPS data and E-PRTR data.

E-PRTR reporting includes ammonia emissions from a large number of agricultural operators. In the Finnish inventory, these facility reported emissions are not taken into account in the inventory reporting (and hence LPS reporting), since all the ammonia emissions in the inventory are calculated in a separate calculation model for agricultural emissions.

Note that the inventory is built up from boiler/process level data and default LPS emissions are calculated as part of the inventory, however, replaced by the reported data in the YLVA system, whenever these data are available and their correctness is checked.

The incorrect coordinates detected in the 2020 NECD review have been corrected to the latest LPS data sets.

Some emissions reported by the facilities for 2020 in the European Environment Agency's (EEA) PRTR database (Industrial Reporting database v5, December 2021) are absent from the LPS data for 2020 set to be reported by 1st May 2022. These include:

- NO_x emissions reported by E-PRTR National IDs 3238 and 11179, SO_x emissions reported by the E-PRTR National ID 100239774 and PM₁₀ emissions reported by E-PRTR National ID 1913. In these cases the 2020 emissions are below the reporting threshold from LPS when the emissions are split between the different stack heights in the LPS reporting.
- NO_x emissions reported by E-PRTR National ID 100131111. The emission data in the reported in the PRTR database for this facility includes emissions also from the fuel use in off-road

working machines. In the air pollutant emission inventory, all emissions from off-road working machines are calculated in the LIPASTO (explained in a more detailed level in the IIR/Part 3 – Transport) calculation system. To avoid double counting, none of the emissions of working machines reported by the facilities themselves are taken into account.

- PCDD/F emissions reported by E-PRTR National ID 1172. These emissions are incorrect in the PRTR database. The emissions are 1000 times too high in the database. The emissions have been corrected to the air pollutant emission inventory and the correct emissions are below the reporting threshold from LPS.
- PCDD/F emissions reported by E-PRTR National ID 3653. The emissions reported by the facility for the first time for 2020 in the PRTR database will be investigated thoroughly with the facility during 2022. Based on this work the possible inclusion of these emissions will be decided for submission 2023.

Data on Finnish LPSs has been submitted annually under the CLRTAP and since 2002 under the NECD.

Uncertainty and time series' consistency

No separate evaluation of uncertainty has been done for LPS data. However, all LPS data are included in the inventory data, for which an uncertainty analysis has been carried out.

Source-specific QA/QC and verification

Normal statistical quality checking related to assessment of magnitude and trends has been carried out.

Source-specific recalculations including changes made in response to the review process

2012

- The definition of the set of Finnish Large Point Sources (LPS) was revised to correspond to the definition of E-PRTR installations. as defined in the revised UNECE Reporting Guidelines (ECE/EB.AIR/97).
- During the preparation of the 2012 submission, it was observed that the conversion of nationally used coordinates into the coordinates in the CLRTAP reporting did not work as believed. A new method to convert the coordinates was introduced.

2017-2018

- The geographical coordinates used in national reporting for point sources is EUREF-FIN and there was a need to carry out a conversion between the level and geographical coordinates. The additional functionality challenges were resolved to the 2019 submission.

2021

- Rewritten documentation
- Correction of some coordinates
- VAHTI-based coordinates updated to more accurate YLVA coordinates.
- Facility locations unified with gridded data.

2022

- Inclusion of a description of the differences between the latest E-PRTR and LPS reporting data sets.

Source-specific planned improvements

None.

8.4 MEMO ITEMS

Changes in chapter	
Update of text	March 2020 KS
Change in methodology	
Other (e.g. language, layout)	

Overall description and methodologies

1 A 3 ai(ii) International aviation cruise

See IIR Part 2 Energy under Aviation.

1 A 3 aii(ii) Domestic aviation cruise

See IIR Part 2 Energy under Aviation.

1 A 3 dii(i) International maritime navigation

See IIR Part 2 Energy under Navigation.

1 A 5 c Multilateral operations

Included under 1A5 (IE)

1 A 3 Transport (fuel used)

.Not applicable (NA). The inventory is based on fuels sold.

6 B Other not included in national total of the entire territory

Not occurring (NO) in Finland.

11 A Volcanoes

There are no volcanoes in Finland.

11 B Forest fires

Not estimated (NE).

11 C Other natural emissions

Not applicable (NA).

REFERENCES

- Aakkula, J., Asikainen, A., Kohl, J., Lehtonen, A., Lehtonen, H., Ollila, P., Regina, K., Salminen, O., Sievänen, R. & Tarja Tuomainen (2019). Maatalous- ja LULUCF-sektorien päästö- ja nielukehitys vuoteen 2050. Valtioneuvoston selvitys- ja tutkimustoiminnan julkaisusarja 20/2019. 70 p. An English abstract: "Development of emissions and sinks in the agricultural and LULUCF sectors until 2050" <http://urn.fi/URN:ISBN:978-952-287-650-8>
- Aasestad K. (2013) Emissions of black carbon and organic carbon in Norway 1990-2011.
- AEAT (2000). UK Fine Particulate Emissions from Industrial Processes. May 2000.
- Aittola, J-P., Paasivirta, J., Vattulainen, A., Sinkkonen, S., Koistinen, J. and Tarhanen, J. (1996). Formation of chloroaromatics at a metal reclamation plant and efficiency of stack filter in their removal from emissions. *Chemosphere* 32. 99-108.
- Anon (2020). Biokaasuohjelmaa valmisteleavan työryhmän loppuraportti (Final report of the biogas working group). Ministry of Economic Affairs and Employment, Ministry of Agriculture and Forestry, Ministry of Transport and Communications, Ministry of the Environment, and Ministry of Finance. Publications of the Ministry of Economic Affairs and Employment 2020:3
- Arnold, M., Kuusisto, S. and Mroueh, U.-M. (1998). Emissions from volatile organic compounds (VOC) in 1996. VTT Publications 1921. Technical Research Centre of Finland. 35 p. (In Finnish).
- Bailey, R.E. (2001). Global hexachlorobenzene emissions. *Chemosphere* 42 167-182.
- BiPRO (2006). Identification, assessment and prioritisation of EU measures to reduce releases of unintentionally produced/released Persistent Organic Pollutants. Interim Report. reference: ENV.D.4/ETU/2005/0068r. 99 p.
- Blomberg, T. (2008). Recalculation of NMVOC emissions from road paving.
- Brenback, M. (2003). Main nutrients in fertilizers 2001/2002. *Vihreä kirja*. Kemira (In Finnish)
- Björndal, H. (1996). Alternatives to persistent organic pollutants. Rapport från kemikalieinspektionen. 4/96. SEPA (Swedish Environment Protection Agency). (In Swedish)
- Best Available Techniques (BAT) Reference Document for the Manufacture of Glass Industrial Emissions Directive 2010/75/EU (Integrated Pollution Prevention and Control)
- CLRTAP/EEA (2006). The Individual review of the CLRTAP air emission inventory of Finland submitted in 2005 (21.12.2006). 13 pages.
- EEA (2013). EMEP/EEA air pollutant emission inventory guidebook - 2013. Technical guidance to prepare national emission inventories. EEA Technical report No 12/2013. European Environmental Agency ISSN 1725-2237. ISBN 978-92-9213-403-7. URL: <http://www.eea.europa.eu/publications/emep-eea-guidebook-2013>

EEA (2009). EMEP/EEA air pollutant emission inventory guidebook - 2009. Technical guidance to prepare national emission inventories. EEA Technical report No 9/2009. European Environment Agency ISBN 978-92-9213-034-3. ISSN 1725-2237. URL:

<http://www.eea.europa.eu/publications/emep-eea-emission-inventory-guidebook-2009>

EEA (2007). EMEP/CORINAIR Emission Inventory Guidebook - 3rd edition. Updated files from December 2007. European Environment Agency Technical report No 30. URL:

<http://www.eea.europa.eu/publications/EMEPCORINAIR5>

EEA (2005). EMEP/CORINAIR Emission Inventory Guidebook - 2005. Sources of HCB emissions. http://reports.eea.europa.eu/EMEPCORINAIR4/en/sources_of_HCB.pdf

EIPPCB (2000). IPPC BAT Reference Document on Iron and Steel Industries (BREF). European Integrated Pollution Prevention and Control Bureau. Seville. March 2000.

Environment Canada (2006). HCB Canadian Inventory for Long Range Transport Modeling. Presentation 17 May 2006 GLBTS Meeting.

FINAVIA (2009). Calculation of international aviation emissions (Ms Niina Rusko).

Finergy (2008). Information on storing of coal. Finnish Energy Industries Federation. Personal communication.

Finland's 6th National Communication to the UNFCCC (2013)

Finlands National Inventory Report to the UNFCCC (2013)

Finnish Glas Industry BAT Group (2001). Conclusions of national BAT group on glass manufacturing (not published)

Finnish Environment Institute (2000) Calculation model for Finnish NMVOC emissions. 32 p.

Finnish Environment Institute (2001). Documentation of the expert estimation for the emission factors of industrial and medical waste incineration. (In Finnish)

Finnish Environment Institute (2007). Waste statistics. Environmental Management Division. Finnish Environment Institute.

Finnish Environment Institute (2004) Ammonia emissions in the UNECE CLRTAP inventory. May 2004. 17 pages

Finnish Environment Institute (2004) Development of the Finnish UNFCCC and UNECE CLRTAP inventories in the agriculture source sector. 22.10.2003. 21 pages

Finnish Environment Institute (2004) NMVOC emissions in the UNFCCC and CLRTAP inventories. 4.2.2004. 29 p.

Finnish Environment Institute (2004) Particle emissions in the UNECE CLRTAP inventory. April 2004. 24 p.

Finnish Environment Institute and Energy Producers (2005). Production of Emission Data – Energy Industries. 103 p. <http://www.ymparisto.fi/download.asp?contentid=46599&lan=fi>

Finnish Environment Institute (2006). Report on Finnish HCB air emissions 1990 – 2004 to the Secretariat of the UNECE Convention on Long-Range Transboundary Air Pollution. 18 December 2006

Finnish Regional Emission Scenario FRES <https://www.sciencedirect.com/science/article/pii/S1352231021005343>

FFIF (1996). The Finnish Forest Industries Federation Annual Report. August 1996.

FFIF (2003). The Finnish Forest Industries Federation Annual Report 2002. Sawmills and board production.

Finnish Biocycle and Biogas Association (2022). Statistics [Tilastot]. In Finnish. Retrieved February 8, 2022, from <https://biokierto.fi/tilastot/>

Finnish Congregations (2013). Number of dead and cremated.

Finnish Fur Sales (2013). Statistics for fur production.

Finnish Petroleum Federation (2014). Oil product and natural gas sales in Finland 2013. http://www.oil.fi/sites/default/files/sivut/sisaltosivu/liitetiedostot/3.4_sales.pdf

Finstad. A., Haakonsen. G and K. Rypdal (2003) Utslipp til luft av partikler I Norge. Dokumentasjon av metode og resultater. Rapportter 2003/15. Statistisk sentralbyrå 2003

Fortum Oil and Gas Ltd. 2002. Revised inventory of Finnish NMVOC emissions in 1988-2001. In Finnish (confidential)

Grönfors. K. (2012). Statistics Finland. Personal communications

Grönroos. J., Nikander. A., Syri. S., Rekolainen. S. and Ekqvist. M. (1998). Agricultural ammonia emissions in Finland. Finnish Environment Institute. The Finnish Environment 206. 65 p. (In Finnish).

Grönroos. J., Mattila. P., Regina. K., Nousiainen. J., Perälä. P., Saarinen. K. and Mikkola-Pusa. J. (2009) Development of ammonia emission inventory in Finland: Description of the revised model and results. Publication available at www.environment.fi/publications>The Finnish Environment.

Hupa. M., Boström. S. and Nermes. M. (1988) Total emissions from energy production in Finland. Insinööritoimisto Prosessikemia Ky. Ministry of Trade and Industry. Energy Department. Series D:162. 62 p. (In Finnish)

Huttula. P. (2002). Written information on emissions from service stations and storage tanks outside the refineries for 2000-2001. Finnish Oil and Gas Federation. (In Finnish)"

Huttunen M. and Kuittinen V. (2017). Finnish national biogas statistics. Data year 2016. Publications of the University of Eastern Finland. Reports and Studies in Forestry and Natural Sciences, 29, University of Eastern Finland, ref.

Information Centre of the Ministry of Agriculture and Forestry (2012) Storage and handling of grain.

Isännäinen. S. (1994). Utilisation of wastewater sludge. Jätevesilietteistä ja niiden hyötykäytöstä. In: Seminaariesitelmiä: Vesiensuojelu. Ympäristönsuojelutekniikan julkaisuja 1994(4): 19–39. Helsinki University of Technology. Laboratory of Environmental Pollution Prevention Technology. (In Finnish).

Joas. A. (2006). Working paper in preparation of EU Implementation plan on POPs under the Stockholm Convention. Annex 1.

Jones. K. (2005). Hexachlorobenzene – Sources, environmental fate and risk characterization. Science Dossier. Euro Chlor. Joas. A. (2006)

Huttunen. M.J. and Kuittinen. V.. Suomen biokaasulaitosrekisteri n:o 17. Publications of the University of Eastern Finland. Reports and s_tureid in Forestry and Natural Sciences No 19 (2014) (In Finnish).

Karvosenoja. N., Johansson. M. and Kupiainen. K. (2002) The importance of primary particulate emissions from non-combustion sources in Finland. Proceedings of 16th International Clean Air and Environment Conference. 19-22.8.2003. Christ Church. New Zealand. Clean Air Society of Austria and New Zealand.. p. 393-398

Karvosenoja. N. (2008) Emission scenario model for regional air pollution. Monographs of the Boreal Environment Research no. 32. p.55.URN:ISBN:978-952-11-3185-1. ISBN 978-952-11-3185-1 (PDF). The publication is available also in printed form ISBN 978-952-11-3184-4.

Kemia-Kemi (2012). Activity data for cement production.

Keränen. S. and Niskanen. R. (1987). Typpilannoituksen vaikutus happamoitumiseen Suomessa. The acidification impact of nitrogen fertilisation in Finland. Helsinki. Ministry of the Environment (HAPRO). 64 p. (In Finnish).

Koljonen, T., Aakkula, J., Honkatukia, J., Soimakallio, S., Haakana, M., Hirvelä, H., Kilpeläinen, H., Kärkkäinen, L., Laitila, J., Lehtilä, A., Lehtonen, H., Maanavilja, L., Ollila, P., Siikavirta, H. & Tuomainen, T. 2020. Hiilineutraali Suomi 2035 - Skenaariot ja vaikutusarviot. 150 s. VTT Technical Research Centre of Finland. VTT Technology, No. 366. An English abstract: "Carbon neutral Finland 2035 – Scenarios and impact assessments". <https://doi.org/10.32040/2242-122X.2020.T366>

KTM (1988). Ministry of Trade and Industry. National total emissions from energy production. Energian-tuotannon kokonaispäästöt Suomessa. KTM. Energiaosasto. D:162.1988 (In Finnish)

Lamberg Heikki (2018) Personal communication Heikki Lamberg. University of Eastern Finland. August 2018.

Lehtonen, H. & Rankinen, K. 2015. Impacts of agri-environmental policy on land use and nitrogen leaching in Finland. Environmental Science and Policy 50: 130–144. doi:10.1016/j.envsci.2015.02.001

Lehtonen, H. & Niskanen, O. 2016. Promoting clover-grass: Implications for agricultural land use in Finland. Land Use Policy 59: 310-319. DOI:10.1016/j.landusepol.2016.09.005

Lehtonen, H., & Niemi, J. 2018. Effects of reducing EU agricultural support payments on production and farm income in Finland. Agricultural and Food Science, 27(2), 124–137. <https://doi.org/10.23986/afsci.67673>

Lehtonen, H. & Rämö, J. 2020. Finland. Section 5.8 (p. 292-318) in: FABLE (2020). Pathways to Sustainable Land-Use and Food Systems. 2020 Report of the FABLE Consortium. Laxenburg and Paris: International Institute for Applied Systems Analysis (IIASA) and Sustainable Development Solutions Network (SDSN). <https://doi.org/10.22022/ESM/12-2020.16896>

LIPASTO (2015). Calculation system for traffic exhaust emissions and energy consumption in Finland. <http://lipasto.vtt.fi/index.htm>

Louhelainen. K.. Vilhunen. P.. Kangas. J. and Terho E.O. (1987) Dust exposure in piggeries. European Journal of Respiratory Diseases. Vol. 71. p. 80-90

Melanen. M.. Ekqvist. M.. Mukherjee. A.B.. Aunela-Tapola. L.. Verta. M. and Salmikangas. T. (1999). Heavy metal emissions in Finland in the 1990's. Suomen ympäristö 329. 92 p. (In Finnish)

MEE. 2012. Study on energy use of waste and emission trading in Finland. Ministry of Employment and Economy. Pöyry Management Consulting. (In Finnish)

MET (2013). Activity data for production rates. Federation of Finnish Metal. Engineering and Electrical Industries. URL: <http://www.met.fi/>

Miljöstyrelsen (2000). Substance Flow Analysis for Dioxins in Denmark. Environmental Project no 570 2000. Miljöstyrelsen

Ministry of the Environment (2002). Air Pollution Control Programme 2010 . The Finnish National Programme for the implementation of Directive 2001/81/EC. approved by the Government on September 26. 2002. 39 p. Available electronically (in Finnish) at <http://www.environment.fi>. please search the site using keyword SY588

Monni. S. & Syri. S.. Savolainen. I. (2003). Uncertainties in the Finnish greenhouse gas emission inventory. Environmental Science & Policy. Vol. 7 (2004) NO:2. 87-98.

Monni. S (2004). Uncertainties in the Finnish 2002 Greenhouse Gas Emission Inventory. VTT. Espoo. 31 p. + app. 18 p. VTT Working Papers 5. <http://www.vtt.fi/inf/pdf/workingpapers/2004/W5.pdf>

Mäkelä. K.. Laurikko. J. and Kanner. H. (2002). Road traffic exhaust gas emissions in Finland. LIISA 2001.1 calculation model. Technical Research Centre of Finland. VTT Research Notes 2177. (In Finnish).

Mäkelä. K.. Laurikko. J. and Kanner. H. (2003). Road traffic exhaust gas emissions in Finland. LIISA 2002 calculation model. Technical Research Centre of Finland. VTT Research Notes 1377. (In Finnish)

Mäkelä. K. (2012). VTT. Gasoline evaporation from vehicles. Personal communication.

Norwegian Emission Inventory (2005). Documentation of methodologies for estimating emissions of greenhouse gases and long-range transboundary air pollutants. Rapport 2005/28. ISBN 82-537-6860-5. 159 p.

NPI (1999). NPI Industry Handbooks. National Pollutant Inventory. Environment Australia. November 1999. <http://www.npi.gov.au/>

Nuutinen. J.. Yli-Pirilä. P.. Hytönen. K. and Kärtevä. J. (2007) Turvetuotannon pöly- ja melupäästöt sekä vaikutukset lähialueen ilmanlaatuun (only in Finnish). Association of Finnish Peat Industries 2007

OSPAR (1999). Oslo and Paris Convention for the protection of the marine environment of the North -East Atlantic. Draft Harp-Haz Guidance on Quantification and Reporting of Polycyclic Aromatic Hydrocarbons (PAH).

Pacyna. J.M... Breivik. K.. Münch. J. and Fudala J. (2003). European atmospheric emissions of selected persistent organic pollutants. 1970-1995. Atmospheric Environment 37

Paulig Ltd (2005) Environmental report 2004.

Pellikka T. (2019) Personal communication Tuula Pellikka VTT. February 2019

Pipatti. R.. Tuhkanen. S.. Mälkiä. P. and Pietilä. R. (2000). Agricultural greenhouse gas emissions and abatement options and costs in Finland. VTT Publications 841. 72 p. Technical Research Centre of Finland. (In Finnish).

Pohjola. V.. Hahkala. M. and Häsänen. E. (1983). Report on emissions from coal. peat and oil combustion processes. VTT Technical Research Centre in Finland. Research Notes 231. Espoo 1983. 139 p. (In Finnish)

Posch. M. (2006) A programme for displaying data on European maps – User Manual.

Rekolainen. S.. Pitkänen. H.. Bleeker. A. and Felix. S. (1995). Nitrogen and phosphorus fluxes from Finnish agricultural areas to the Baltic Sea. Nordic Hydrology 26: 55 B 72.

RT. (2013). Information on production rates. Confederation of Finnish Construction Industries. Information on the website URL: <http://www.rakennusteollisuusrt.fi/> and personal communication.

Ruuskanen. J. (2000). Environmental protection technology for reducing dioxine emissions in the 1990's. Dioksiineja vähentävät ympäristönsuojelukeinot 1990-luvulla ja uudet mahdollisuudet. Ympäristö- ja terveystiedote 3:2000. (In Finnish)

Saarinen. K.. Lammi. R.. Silvo. K. and Hietämäki. M. (2004) Emission data production – Forest Industries. 73 p. Website at www.ymparisto.fi > Yritykset ja yhteisöt > Päästöt > Päästörekisterit > Päästötiedon tuottaminen > Aineistoa päästöjen määrittämisen tueksi (In Finnish)

Saarinen. K (ed (2004) Emission data production – Energy Industries. Website at www.ymparisto.fi > Yritykset ja yhteisöt > Päästöt > Päästörekisterit > Päästötiedon tuottaminen > Aineistoa päästöjen määrittämisen tueksi (In Finnish)

SEPA1996. Alternatives to persistent organic pollutants. Rapport från kemikalieinspektionen. 4/96. Swedish Environment Protection Agency. SEPA.

Seppänen. H. & Matinlassi. T. (1998). Environmental care programs at Finnish farms 1995 B 1997. Maaseutukeskusten liitto (Rural Advisory Centres). 19 p. (In Finnish).

Statistics Finland (2014). Preliminary information on 2012 data from Statistics Finland.

Statistics Finland (1995). Wastes from Manufacturing and Related Industries 1992. Official Statistics of Finland. Environment 1995 (7). 162 p.

Statistic Finland (2009) Greenhouse gas emissions in Finland 1990-2007; National Inventory Report to the European Union

Suoheimo. P. Grönroos. J.. Karvosenoja. N.. Petäjä. J.. Saarinen. K.. Savolahti. M.. Silvo. K (2015). Impacts of the implementation of the Revision of National Emission Ceilings Directive and the Proposed Medium Combustion Plants Directive in Finland. Reports of the Finnish Environment Institute 6/2015. https://helda.helsinki.fi/bitstream/handle/10138/153981/SYKEra_6_2015.pdf?sequence=3 78 p. In Finnish.

SYKE. (2001). Expert estimation based on UNEP 1999.

SYKE. (2007a). Expert estimation based on Aittola (1996) and Joas (2006).

Takai. H.. Pedersen. S.. Johnsen. O.. Metz. J. H.M.. Groot Koerkamp. P.W.G.. Uenk. G.H.. Phillips. V.R.. Holden. M.R.. Sneath. R.W.. Short. J.L.. White. R.P.. Hartung. J.. Seedorf. J.. Schröder. M.. Linkert. K.H. and Wathes. C.M. (1998) Concentrations and emissions of airborne dust in livestock buildings in Northern Europe. Journal of Agricultural Engineering Research. Vol 70. p. 59-77

The Federation of Finnish Technology Industries (2014) Activity data for production rates. Technology Industries of Finland. URL: <http://www.teknologiateollisuus.fi>

TIKE (2014) Yearbook of Farm Statistics 2013. Information Center of the Ministry of Agriculture and Forestry. URL: <http://tike.mmm.fi/>

TNO (2002). The Co-ordinated European Programme on Particulate Matter Emission Inventories. Projections and Guidance (CEPMEIP). <http://www.air.sk/tno/cepmeip/>

TNO (1995). TNO-Report TNO-MEP - 95/247: Technical paper to the OSPARCOM-HELCOM-UNECE emission Inventory of Heavy Metals and POP.

Toda. E. (2005). POPs and heavy metals emission inventory of Japan. TFEIP & ES-PREME Workshop on "Heavy Metals and POPs Emissions. Inventories and Projections". 18-19 October 2005. Rovaniemi. Finland.

Tsupari. E.. Tormonen. K.. Monni. S.. Vahlman. T.. Kolsi. A. and Linna. V. (2006). Emission factors for nitrous oxide (N₂O) and methane (CH₄) from Finnish power and heating plants and small-scale combustion. VTT. Espoo. 94 s. + liitt. 7 s. VTT Working Papers : 43
ISBN 951-38-6595-9

UBA (1998). Umwelt Bundesamt. Investigation of emissions and abatement measures for persistent organic pollutants in the FRG. research report 295 44 365. UBA-*FB 98-115/e. 75/98

UNEP (1999). United Nations Environment Programme. Dioxin and furan inventories. National and regional emissions of PCDD/PCDF. May 1999

UNEP (2005) Standardized Toolkit for Identification and Quantification of Dioxin and Furan Releases. 2nd edition. February 2005

USEPA (1997). United States Environmental Protection Agency. Office of Air Quality Planning And Standards. Locating and Estimating Air Emissions from Sources of Dioxins and Furans. USEPA. May 1997.

USEPA (1998). Locating and Estimating Air Emissions from Sources of Polycyclic Organic Matter (EPA-454/R-98-014) July 1998

Westerlund.. K-G. (2001) Metal emissions from Stockholm Traffics Wear of Brake Linings; Reports from SLB-analys. 2:2001; Environment and Health Protection Administration in Stockholm: Stockholm 2001